

Universal Deformations and Inhomogeneities in Isotropic Cauchy Elasticity*

Arash Yavari^{†1,2}

¹*School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

²*The George W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

June 5, 2024

Dedicated to Professor Kaushik Bhattacharya on the occasion of his 60th birthday.

Abstract

For a given class of materials, *universal deformations* are those deformations that can be maintained in the absence of body forces and by applying solely boundary tractions. For inhomogeneous bodies, in addition to the universality constraints that determine the universal deformations, there are extra constraints on the form of the material inhomogeneities—*universal inhomogeneity constraints*. Those inhomogeneities compatible with the universal inhomogeneity constraints are called *universal inhomogeneities*. In a Cauchy elastic solid, stress at a given point and at an instance of time is a function of strain at that point and that exact moment in time, without any dependence on prior history. A Cauchy elastic solid does not necessarily have an energy function, i.e., Cauchy elastic solids are, in general, non-hyperelastic (or non-Green elastic). In this paper we characterize universal deformations in both compressible and incompressible inhomogeneous isotropic Cauchy elasticity. As Cauchy elasticity includes hyperelasticity, one expects the universal deformations of Cauchy elasticity to be a subset of those of hyperelasticity both in the compressible and incompressible cases. It is also expected that the universal inhomogeneity constraints to be more stringent than those of hyperelasticity, and hence, the set of universal inhomogeneities to be smaller than that of hyperelasticity. We prove the somewhat unexpected result that the sets of universal deformations of isotropic Cauchy elasticity and isotropic hyperelasticity are identical, in both the compressible and incompressible cases. We also prove that their corresponding universal inhomogeneities are identical as well.

Keywords: Universal deformation, universal inhomogeneity, Cauchy elasticity, hyperelasticity, Green elasticity, isotropic solids.

1 Introduction

Within a given class of materials, universal deformations refer to those deformations that can be maintained in the absence of body forces and by applying only boundary tractions, for any member of the material class. Universal deformations do not depend on the particular material within the class. However, the boundary tractions necessary to sustain a universal deformation depend on the specific material. Universal deformations have played a crucial role in nonlinear elasticity and anelasticity (in the sense of [Eckart \[1948\]](#)): i) They have had an important organizational role in the semi-inverse solutions in nonlinear elasticity [[Knowles, 1979](#), [Polignone and Horgan, 1991](#), [De Pascalis et al., 2009](#), [Tadmor et al., 2012](#), [Goriely, 2017](#)], and more recently in anelasticity [[Kumar and Yavari, 2023](#)] and viscoelasticity [[Sadik and Yavari, 2024](#)]. ii) They offer

*To appear in the *Proceedings of the Royal Society A*.

[†]Corresponding author, e-mail: arash.yavari@ce.gatech.edu

guidance for designing experiments for determining the constitutive relations of a specific material [Rivlin and Saunders, 1951, Saccomandi, 2001].¹ iii) All the existing exact solutions of defects in nonlinear solids are related to universal deformations [Wesolowski and Seeger, 1968, Gairola, 1979, Zubov, 1997, Yavari and Goriely, 2012a,b, 2013a, 2014, Golgoon and Yavari, 2018b]. iv) Universal deformations have been important for finding exact solutions for distributed finite eigenstrains in nonlinear solids and the nonlinear analogues of Eshelby’s inclusion problem [Yavari and Goriely, 2013b, 2015, Golgoon and Yavari, 2018a, Yavari, 2021b]. v) These exact solutions have been used as benchmark problems in computational mechanics [Dragoni, 1996, Saccomandi, 2001, Chi et al., 2015, Shojaei and Yavari, 2018]. vi) Universal deformations have been used in deriving effective properties for nonlinear composites [Hashin, 1985, Lopez-Pamies et al., 2012, Golgoon and Yavari, 2021].

The systematic study of universal deformations began in the 1950s by Jerry Ericksen [Ericksen, 1954, 1955] for homogeneous compressible and incompressible isotropic solids. His work was influenced by the earlier contributions of Ronald Rivlin [Rivlin, 1948, 1949a,b]. Ericksen [1955] proved that in homogeneous compressible isotropic solids, universal deformations are homogeneous. The characterization of universal deformations in the presence of internal constraints is a particularly challenging problem [Saccomandi, 2001]. Ericksen [1954] discovered four families of universal deformations for incompressible isotropic elastic solids. Initially, he speculated that a deformation with constant principal invariants is homogeneous, but this conjecture was proven incorrect [Fosdick, 1966]. Subsequently, a fifth family of universal deformations was found [Singh and Pipkin, 1965, Klingbeil and Shield, 1966]. The fifth family of universal deformations have constant principal invariants but are not homogeneous. As of now, it is not known whether there exist additional inhomogeneous constant-principal invariant universal deformations (Ericksen’s problem). The following are the six known families of universal deformations:

- Family 0: Homogeneous deformations
- Family 1: Bending, stretching, and shearing of a rectangular block
- Family 2: Straightening, stretching, and shearing of a sector of a cylindrical shell
- Family 3: Inflation, bending, torsion, extension, and shearing of a sector of an annular wedge
- Family 4: Inflation/inversion of a sector of a spherical shell
- Family 5: Inflation, bending, extension, and azimuthal shearing of an annular wedge

There have been several attempts to solve Ericksen’s problem in the past few decades [Marris and Shiau, 1970, Kafadar, 1972, Marris, 1975, 1982]. Fosdick and Schuler [1969] showed that for the case of plane deformations with uniform transverse stretch there are no new solutions other than the known families. Fosdick [1971] reached the same conclusion for radially-symmetric deformations.

In a *simple material*, stress at any given point depends only on the history of the deformation gradient at that point up to time t [Noll, 1958]. Carroll [1967] demonstrated that the above known universal deformations of homogenous incompressible isotropic elastic solids are universal for simple materials as well.

The study of universal deformations was extended to inhomogeneous anisotropic solids in [Yavari and Goriely, 2021, Yavari, 2021a, Yavari and Goriely, 2023].² These comprehensive studies included both compressible and incompressible isotropic, transversely isotropic, orthotropic, and monoclinic solids. For these three classes of compressible anisotropic solids, it was shown that universal deformations are homogenous, and the material preferred directions are uniform. Additionally, for isotropic solids and each of the three classes of anisotropic solids, the corresponding universal inhomogeneities—these represent inhomogeneities in the energy function that are compatible with the universality constraints—were characterized. The corresponding universal inhomogeneities for each of the above six known families of universal deformations were determined for inhomogeneous isotropic and the three classes of inhomogeneous incompressible anisotropic solids.

In linear elasticity, universal displacements are the counterparts to universal deformations [Truesdell, 1966, Gurtin, 1972, Carroll, 1973, Yavari et al., 2020]. Yavari et al. [2020] demonstrated the explicit de-

¹The following quote from [Doyle and Ericksen, 1956, p. 89] explains it best: “From the standpoint of attempting to determine the form of Σ for a given material by comparing general solutions with results of experiment, it appears that the solutions which are most useful are those which correspond to deformations which can be produced in every material of the type considered by the application of surface tractions only.” Here, Σ is the energy function and by “general” they mean “universal”.

²Prior to our work, there had been some limited work on universal deformations in anisotropic solids [Ericksen and Rivlin, 1954].

pendence of universal displacements on the symmetry class of the material. Specifically, the larger the symmetry group, the larger the corresponding set of universal displacements. Therefore, isotropic solids have the largest set of universal displacements, while triclinic solids possess the smallest set of universal displacements. The investigation into universal displacements has also been extended to inhomogeneous solids [Yavari and Goriely, 2022b] and linear anelasticity [Yavari and Goriely, 2022a].

A class of materials with internal constraints, significant in engineering applications, consists of materials reinforced with inextensible fibers [Adkins and Rivlin, 1955, Pipkin and Rogers, 1971, Pipkin, 1974, 1979]. Despite their importance, there is a scarcity of literature on the universal deformations of fiber-reinforced solids. Beskos [1972] studied homogeneous compressible isotropic solids reinforced with inextensible fibers, investigating whether universal deformations of incompressible isotropic solids are universal for this class as well. Families 1, 2, 3, and 4 were specifically analyzed, revealing that certain subsets of these families are universal for specific fiber distributions. Interestingly, all these universal deformations are found to be homogeneous except for the shearing of a circular tube with circumferential fibers. Beatty [1978] further examined homogeneous compressible isotropic solids reinforced by a single family of inextensible fibers. He investigated the problem of finding all those fiber distributions for which homogenous deformations are universal. He proved that there are only three types of such fiber distributions, all characterized by straight fibers. In a recent study, Yavari [2023] studied universal displacements in compressible anisotropic linear elastic solids reinforced by a single family of inextensible fibers. For each symmetry class, and under the assumption of a uniform distribution of straight fibers respecting the corresponding symmetry, the respective universal displacements were characterized. Interestingly, it was observed that, except for triclinic and cubic solids, the presence of inextensible fibers enlarges the set of universal displacements within the other five classes.³

In recent years, Ericksen’s analysis has been extended to anelasticity. Yavari and Goriely [2016] showed that in compressible anelasticity, universal deformations are covariantly homogeneous. Universal deformations and eigestrains in incompressible anelasticity were studied by Goodbrake et al. [2020]. It was observed that the six known families of universal deformations are invariant under specific Lie subgroups of the special Euclidean group. There are also some recent studies of universal deformations and eigenstrains in accreting bodies [Yavari and Pradhan, 2022, Yavari et al., 2023, Pradhan and Yavari, 2023]. There have also been studies of universal deformations in liquid crystal elastomers [Lee and Bhattacharya, 2023, Mihai and Goriely, 2023].

In this paper, we extend the study of universal deformations and inhomogeneities to inhomogeneous compressible and incompressible isotropic Cauchy elasticity. Cauchy elastic solids may not necessarily have an energy function and include hyperelastic solids (Green elastic solids) as a special case. This suggests that Cauchy elasticity may have more stringent universality and universal inhomogeneity constraints compared to those of hyperelasticity. This, in turn, leads one to anticipate smaller sets of universal deformations and universal inhomogeneities for Cauchy elasticity compared to those of hyperelasticity. We prove the somewhat unexpected result that the universal deformations and inhomogeneities of Cauchy elasticity are identical to those of Green elasticity in both the compressible and incompressible cases.

This paper is organized as follows. In §2, Cauchy elasticity is briefly reviewed. Universal deformations of inhomogeneous compressible isotropic Cauchy elasticity are characterized in §3. In §4, the same problem is studied for inhomogeneous incompressible isotropic Cauchy elasticity. Conclusions are given in §5.

2 Cauchy Elasticity

Let us consider a body that in its undeformed configuration is identified with an embedded submanifold \mathcal{B} of the Euclidean ambient space \mathcal{S} . The flat metric of the Euclidean ambient space is denoted by \mathbf{g} and the induced metric on the body in its reference configuration by $\mathbf{G} = \mathbf{g}|_{\mathcal{B}}$. Deformation is a map from \mathcal{B} to the ambient space, i.e., $\varphi : \mathcal{B} \rightarrow \mathcal{C} \subset \mathcal{S}$, where $\mathcal{C} = \varphi(\mathcal{B})$ is the current configuration. The tangent map of φ is the so-called deformation gradient $\mathbf{F} = T\varphi$ (a metric independent map), which at each material point $X \in \mathcal{B}$ is a linear map $\mathbf{F}(X) : T_X\mathcal{B} \rightarrow T_{\varphi(X)}\mathcal{C}$. With respect to the coordinate charts $\{X^A\}$ and $\{x^a\}$ for

³It should be noted that a fiber-reinforced solid with straight fibers cannot be isotropic.

\mathcal{B} and \mathcal{C} , respectively, deformation gradient has components $F^a{}_A = \frac{\partial \varphi^a}{\partial X^A}$. The transpose of deformation gradient \mathbf{F}^\top has components $(F^\top)^A{}_a = g_{ab} F^b{}_B G^{AB}$. The right Cauchy-Green strain is defined as $\mathbf{C} = \mathbf{F}^\top \mathbf{F}$ and has components $C^A{}_B = (F^\top)^A{}_a F^a{}_B$. Thus, $C_{AB} = (g_{ab} \circ \varphi) F^a{}_A F^b{}_B$, which means that the right Cauchy-Green strain is the pull-back of the spatial metric to the reference configuration, i.e., $\mathbf{C}^\flat = \varphi^* \mathbf{g}$, where \flat is the flat operator induced by the metric \mathbf{G} (which lowers indices). The left Cauchy-Green strain is defined as $\mathbf{B}^\sharp = \varphi^*(\mathbf{g}^\sharp)$, which has components $B^{AB} = F^{-A}{}_a F^{-B}{}_b g^{ab}$. The spatial analogue of \mathbf{C}^\flat is defined as $\mathbf{c}^\flat = \varphi_* \mathbf{G}$, which has components $c_{ab} = F^A{}_A F^B{}_B G_{AB}$. Similarly, the spatial analogue of \mathbf{B}^\sharp is $\mathbf{b}^\sharp = \varphi_*(\mathbf{G}^\sharp)$, which has components $b^{ab} = F^a{}_A F^b{}_B G^{AB}$. Recall that $\mathbf{b} = \mathbf{c}^{-1}$. The two tensors \mathbf{C} and \mathbf{b} have the same principal invariants I_1 , I_2 , and I_3 , which are defined as [Ogden, 1984, Marsden and Hughes, 1994] $I_1 = \text{tr } \mathbf{b} = b^{ab} g_{ab}$, $I_2 = \frac{1}{2} (I_1^2 - \text{tr } \mathbf{b}^2) = \frac{1}{2} (I_1^2 - b^{ab} b^{cd} g_{ac} g_{bd})$, and $I_3 = \det \mathbf{b}$.

In Cauchy elasticity, stress at a point and a given moment in time is explicitly a function of strain at that point and that particular moment in time [Cauchy, 1828, Truesdell, 1952, Truesdell and Noll, 2004]. However, an energy function does not necessarily exist.⁴ In terms of the first Piola-Kirchhoff stress [Truesdell, 1952, Truesdell and Noll, 2004, Ogden, 1984]

$$\mathbf{P} = \hat{\mathbf{P}}(X, \mathbf{F}, \mathbf{G}, \mathbf{g}). \quad (2.1)$$

One can show that objectivity implies that the second Piola-Kirchhoff stress has the following functional form [Truesdell and Noll, 2004]

$$\mathbf{S} = \hat{\mathbf{S}}(X, \mathbf{C}^\flat, \mathbf{G}). \quad (2.2)$$

For an isotropic solid one has the following classic representation [Rivlin and Ericksen, 1955, Wang, 1969, Boehler, 1977]

$$\mathbf{S} = \Lambda_0 \mathbf{G}^\sharp + \Lambda_1 \mathbf{C}^\sharp + \Lambda_{-1} \mathbf{C}^{-\sharp}, \quad (2.3)$$

where $\Lambda_i = \Lambda_i(X, I_1, I_2, I_3)$, $i = -1, 0, 1$, and \sharp is the sharp operator induced from the metric \mathbf{G} (which raises indices). For incompressible isotropic Cauchy solids $I_3 = 1$, and one has

$$\mathbf{S} = -p \mathbf{C}^{-\sharp} + \Lambda_0 \mathbf{G}^\sharp + \Lambda_1 \mathbf{C}^\sharp, \quad (2.4)$$

where $p = p(X, t)$ is the Lagrange multiplier associated with the incompressibility constraint $J = \sqrt{I_3} = 1$, and $\Lambda_i = \Lambda_i(X, I_1, I_2)$, $i = 0, 1$.

In terms of the Cauchy stress, the constitutive equations of compressible isotropic Cauchy elastic solids are written as

$$\boldsymbol{\sigma} = \alpha \mathbf{g}^\sharp + \beta \mathbf{b}^\sharp + \gamma \mathbf{c}^\sharp, \quad (2.5)$$

where $\alpha = \alpha(X, I_1, I_2, I_3)$, $\beta = \beta(X, I_1, I_2, I_3)$, and $\gamma = \gamma(X, I_1, I_2, I_3)$ are arbitrary response functions. Similarly, for incompressible isotropic Cauchy elastic solids one has

$$\boldsymbol{\sigma} = -p \mathbf{g}^\sharp + \beta \mathbf{b}^\sharp + \gamma \mathbf{c}^\sharp, \quad (2.6)$$

where $\beta = \beta(X, I_1, I_2)$ and $\gamma = \gamma(X, I_1, I_2)$ are arbitrary response functions. In components, they read $\sigma^{ab} = \alpha g^{ab} + \beta b^{ab} + \gamma c^{ab}$ and $\sigma^{ab} = -p g^{ab} + \beta b^{ab} + \gamma c^{ab}$, respectively.

Green and Naghdi [1971] showed that Cauchy elasticity is consistent with the first and second laws of thermodynamics. They demonstrated that over a closed path in the space of strains the net work in a Cauchy elastic solid may not vanish. They explicitly demonstrated this in the case of a linear stress-strain relationship and observed that the lack of major symmetries is responsible for the nonzero work.⁵ There is no consensus in the literature of nonlinear elasticity on Cauchy elasticity being a viable theory; while some dismiss it [Rivlin, 1986, Casey, 2005, Carroll, 2009, Rajagopal, 2011], others appear to accept it [Truesdell, 1952, Ericksen, 1956, Truesdell and Noll, 2004, Ogden, 1984, Kružík and Roubíček, 2019, Bordiga et al.,

⁴It is important to note that Cauchy elasticity does not encompass all elastic solids. In recent years, there has been some interest in implicit constitutive equations, e.g., constitutive equations of the form $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$ [Morgan, 1966, Rajagopal, 2003, 2007]. Cauchy elasticity is a subset of this class of solids.

⁵The same thing was shown fifty years later in [Scheibner et al., 2020] when formulating the so-called “odd” elasticity. These authors clearly were not aware of similar developments in the literature of nonlinear elasticity.

2022]. Our motivation for studying Cauchy elasticity is because of its promise in describing the mechanics of active matter at large strains. Active matter may have access to external sources of energy, and hence, the net work done in a closed loop in the strain space may not vanish. The recent interest in the physics literature in the so-called “odd” elasticity [Scheibner et al., 2020, Fruchart et al., 2023], which is simply linearized non-hyperelastic Cauchy elasticity, is another motivation for revisiting Cauchy elasticity.

3 Universal Deformations in Compressible Isotropic Cauchy Elasticity

Let us consider a compressible isotropic Cauchy elastic body deforming in the Euclidean ambient space. In a local coordinate chart $\{x^a\}$, which may be curvilinear, the Cauchy stress has the following representation

$$\sigma^{ab} = \alpha g^{ab} + \beta b^{ab} + \gamma c^{ab}. \quad (3.1)$$

When there are no body forces present, the equilibrium equations read $\sigma^{ab}|_b = 0$.⁶ Thus⁷

$$\sigma^{ab}|_b = \beta b^{ab}|_b + \gamma c^{ab}|_b + \alpha_{,b} g^{ab} + \beta_{,b} b^{ab} + \gamma_{,b} c^{ab} = 0. \quad (3.2)$$

Note that

$$\begin{aligned} \alpha_{,b} &= F^{-A}{}_b \frac{\partial \alpha}{\partial X^A} + \frac{\partial \alpha}{\partial I_1} I_{1,b} + \frac{\partial \alpha}{\partial I_2} I_{2,b} + \frac{\partial \alpha}{\partial I_3} I_{3,b}, \\ \beta_{,b} &= F^{-A}{}_b \frac{\partial \beta}{\partial X^A} + \frac{\partial \beta}{\partial I_1} I_{1,b} + \frac{\partial \beta}{\partial I_2} I_{2,b} + \frac{\partial \beta}{\partial I_3} I_{3,b}, \\ \gamma_{,b} &= F^{-A}{}_b \frac{\partial \gamma}{\partial X^A} + \frac{\partial \gamma}{\partial I_1} I_{1,b} + \frac{\partial \gamma}{\partial I_2} I_{2,b} + \frac{\partial \gamma}{\partial I_3} I_{3,b}. \end{aligned} \quad (3.3)$$

These can be written more concisely as

$$\begin{aligned} \alpha_{,b} &= F^{-A}{}_b \alpha_{,A} + \alpha_1 I_{1,b} + \alpha_2 I_{2,b} + \alpha_3 I_{3,b}, \\ \beta_{,b} &= F^{-A}{}_b \beta_{,A} + \beta_1 I_{1,b} + \beta_2 I_{2,b} + \beta_3 I_{3,b}, \\ \gamma_{,b} &= F^{-A}{}_b \gamma_{,A} + \gamma_1 I_{1,b} + \gamma_2 I_{2,b} + \gamma_3 I_{3,b}, \end{aligned} \quad (3.4)$$

where

$$\begin{aligned} \alpha_{,A} &= \frac{\partial \alpha}{\partial X^A}, \quad \beta_{,A} = \frac{\partial \beta}{\partial X^A}, \quad \gamma_{,A} = \frac{\partial \gamma}{\partial X^A}, \quad A = 1, 2, 3, \\ \alpha_i &= \frac{\partial \alpha}{\partial I_i}, \quad \beta_i = \frac{\partial \beta}{\partial I_i}, \quad \gamma_i = \frac{\partial \gamma}{\partial I_i}, \quad i = 1, 2, 3. \end{aligned} \quad (3.5)$$

Substituting (3.4) into (3.2) one obtains

$$\begin{aligned} &\beta b^{ab}|_b + \gamma c^{ab}|_b \\ &+ I_{1,b} g^{ab} \alpha_1 + I_{2,b} g^{ab} \alpha_2 + I_{3,b} g^{ab} \alpha_3 \\ &+ I_{1,b} b^{ab} \beta_1 + I_{2,b} b^{ab} \beta_2 + I_{3,b} b^{ab} \beta_3 \\ &+ I_{1,b} c^{ab} \gamma_1 + I_{2,b} c^{ab} \gamma_2 + I_{3,b} c^{ab} \gamma_3 \\ &+ F^{-A}{}_b \delta^{ab} \alpha_{,A} + F^{-A}{}_b b^{ab} \beta_{,A} + F^{-A}{}_b c^{ab} \gamma_{,A} = 0. \end{aligned} \quad (3.6)$$

Note that α , β , and γ are arbitrary functions and their derivatives are independent. Therefore, for the equilibrium equations to hold for any compressible isotropic Cauchy elastic solid, the coefficient of each

⁶ $(.)|_a$ denotes covariant derivative with respect to the vector $\frac{\partial}{\partial x^a}$. In Cartesian coordinates, this reduces to a partial derivative. Also, it should be noted that for any scalar field f , $f|_a = f_{,a}$.

⁷We have used the fact that a Riemannian metric is compatible with its Levi-Civita connection, i.e., $g^{ab}|_c = 0$, and hence $g^{ab}|_b = 0$, where summation over repeated indices is implied.

derivative must vanish. Thus

$$\begin{cases} b^{ab}|_b = c^{ab}|_b = 0, \end{cases} \quad (3.7)$$

$$\begin{cases} g^{ab}I_{1,b} = g^{ab}I_{2,b} = g^{ab}I_{3,b} = 0, \end{cases} \quad (3.8)$$

$$\begin{cases} b^{ab}I_{1,b} = b^{ab}I_{2,b} = b^{ab}I_{3,b} = 0, \end{cases} \quad (3.9)$$

$$\begin{cases} c^{ab}I_{1,b} = c^{ab}I_{2,b} = c^{ab}I_{3,b} = 0, \end{cases} \quad (3.10)$$

$$\begin{cases} F^{-A}_b \delta^{ab} \alpha_{,A} = F^{-A}_b b^{ab} \beta_{,A} = F^{-A}_b c^{ab} \gamma_{,A} = 0. \end{cases} \quad (3.11)$$

The universality constraints (3.8) imply that I_1, I_2, I_3 are constant. Then, the universality constraints (3.9) and (3.10) are trivially satisfied. From (3.11), one concludes that $\alpha_{,A} = \beta_{,A} = \gamma_{,A} = 0$, i.e., the body must be homogeneous. Thus, in summary we have concluded that

$$I_1, I_2, I_3 \text{ are constant, and } b^{ab}|_b = c^{ab}|_b = 0. \quad (3.12)$$

Using these and the compatibility equations, one can show that the universal deformations must be homogeneous [Ericksen, 1955]. Thus, we have proved the following two results.

Proposition 3.1. *The set of universal deformations of homogeneous compressible isotropic Cauchy elastic solids is the set of all homogeneous deformations.*

Proposition 3.2. *Inhomogeneous compressible isotropic Cauchy elastic solids do not admit universal deformations.*

Remark 3.3. We observe that the set of universal deformations of homogeneous compressible isotropic Cauchy elastic solids is identical to that of homogeneous compressible isotropic hyperelastic solids [Ericksen, 1955]. Additionally, it is noteworthy that neither inhomogeneous compressible isotropic Cauchy elastic solids nor inhomogeneous compressible isotropic hyperelastic solids admit universal deformations [Yavari, 2021a].

4 Universal Deformations in Incompressible Isotropic Cauchy Elasticity

For an incompressible isotropic Cauchy elastic solid the Cauchy stress has the following component representation

$$\sigma^{ab} = -p g^{ab} + \beta b^{ab} + \gamma c^{ab}, \quad (4.1)$$

where a curvilinear coordinate chart $\{x^a\}$ is assumed. Equilibrium equations in the absence of body forces read $\sigma^{ab}|_b = \sigma^{ab}_{,b} + \gamma^a_{bc} \sigma^{cb} + \gamma^b_{bc} \sigma^{ac} = 0$, where γ^a_{bc} (not to be confused with the scalar field γ in (4.1)) are the Levi-Civita connection components corresponding to the metric \mathbf{g} . Thus

$$p_{,n} g^{an} = \beta b^{ab}|_n + \gamma c^{ab}|_n + \beta_{,n} b^{ab} + \gamma_{,n} c^{ab}, \quad (4.2)$$

or⁸

$$p_{,a} = \beta b^n_{a|n} + \gamma c^n_{a|n} + \beta_{,n} b^n_a + \gamma_{,n} c^n_a. \quad (4.3)$$

Thus, $dp = p_{,a} dx^a = (\beta b^n_{a|n} + \gamma c^n_{a|n} + \beta_{,n} b^n_a + \gamma_{,n} c^n_a) dx^a = \boldsymbol{\xi}$, where d is the exterior derivative. This means the one-form $\boldsymbol{\xi}$ must be exact. A necessary condition for $\boldsymbol{\xi}$ to be exact is $d\boldsymbol{\xi} = 0$. This implies that $p_{,ab} = p_{,ba}$, which is equivalent to $p|_{ab} = p|_{ba}$. Thus

$$p|_{ab} = \beta b^n_{a|nb} + \gamma c^n_{a|nb} + \beta_{,n} b^n_{a|b} + \gamma_{,n} c^n_{a|b} + \beta_{,b} b^n_{a|n} + \gamma_{,b} c^n_{a|n} + (\beta_{,n})|_b b^n_a + (\gamma_{,n})|_b c^n_a, \quad (4.4)$$

must be symmetric in (a, b) .

⁸Note that $b^n_a = b^{nm} g_{ma}$, and $b_a^n = g_{am} b^{mn}$, which are equal. Thus, we use $b^n_a = b^n_a = b_a^n$. Similarly, the same notation is used for \mathbf{c} .

Note that

$$\beta_{,n} = F^{-A}{}_n \beta_{,A} + \beta_1 I_{1,n} + \beta_2 I_{2,n}, \quad \gamma_{,n} = F^{-A}{}_n \gamma_{,A} + \gamma_1 I_{1,n} + \gamma_2 I_{2,n}, \quad (4.5)$$

where

$$\begin{aligned} \beta_i &= \frac{\partial \beta}{\partial I_i}, \quad \gamma_i = \frac{\partial \gamma}{\partial I_i}, \quad i = 1, 2, \\ \beta_{,A} &= \frac{\partial \beta}{\partial X^A}, \quad \gamma_{,A} = \frac{\partial \gamma}{\partial X^A}, \quad A = 1, 2, 3. \end{aligned} \quad (4.6)$$

Also

$$\begin{aligned} (\beta_{,n})|_b &= \beta_1 I_{1|nb} + \beta_2 I_{2|nb} + \beta_{1,b} I_{1,n} + \beta_{2,b} I_{2,n} + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) \beta_{,A} \\ &\quad + F^{-A}{}_n (F^{-B}{}_b \beta_{,AB} + \beta_{1,A} I_{1,b} + \beta_{2,A} I_{2,b}), \\ &= \beta_1 I_{1|nb} + \beta_2 I_{2|nb} + I_{1,b} I_{1,n} \beta_{11} + I_{2,b} I_{2,n} \beta_{22} + (I_{1,b} I_{2,n} + I_{1,n} I_{2,b}) \beta_{12} \\ &\quad + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) \beta_{,A} + \frac{1}{2} (F^{-A}{}_n F^{-B}{}_b + F^{-B}{}_n F^{-A}{}_b) \beta_{,AB} \\ &\quad + (F^{-A}{}_b I_{1,n} + F^{-A}{}_n I_{1,b}) \beta_{1,A} + (F^{-A}{}_b I_{2,n} + F^{-A}{}_n I_{2,b}) \beta_{2,A}, \end{aligned} \quad (4.7)$$

and

$$\begin{aligned} (\gamma_{,n})|_b &= \gamma_1 I_{1|nb} + \gamma_2 I_{2|nb} + \gamma_{1,b} I_{1,n} + \gamma_{2,b} I_{2,n} + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) \gamma_{,A} \\ &\quad + F^{-A}{}_n (F^{-B}{}_b \gamma_{,AB} + \gamma_{1,A} I_{1,b} + \gamma_{2,A} I_{2,b}), \\ &= \gamma_1 I_{1|nb} + \gamma_2 I_{2|nb} + I_{1,b} I_{1,n} \gamma_{11} + I_{2,b} I_{2,n} \gamma_{22} + (I_{1,b} I_{2,n} + I_{1,n} I_{2,b}) \gamma_{12} \\ &\quad + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) \gamma_{,A} + \frac{1}{2} (F^{-A}{}_n F^{-B}{}_b + F^{-B}{}_n F^{-A}{}_b) \gamma_{,AB} \\ &\quad + (F^{-A}{}_b I_{1,n} + F^{-A}{}_n I_{1,b}) \gamma_{1,A} + (F^{-A}{}_b I_{2,n} + F^{-A}{}_n I_{2,b}) \gamma_{2,A}, \end{aligned} \quad (4.8)$$

where

$$\begin{aligned} \beta_{ij} &= \frac{\partial^2 \beta}{\partial I_i \partial I_j}, \quad \gamma_{ij} = \frac{\partial^2 \gamma}{\partial I_i \partial I_j}, \quad i \leq j = 1, 2, \\ \beta_{i,A} &= \frac{\partial^2 \beta}{\partial I_i \partial X^A}, \quad \gamma_{i,A} = \frac{\partial^2 \gamma}{\partial I_i \partial X^A}, \quad i = 1, 2, \quad A = 1, 2, 3, \\ \beta_{,AB} &= \frac{\partial^2 \beta}{\partial X^A \partial X^B}, \quad \gamma_{,AB} = \frac{\partial^2 \gamma}{\partial X^A \partial X^B}, \quad A \leq B = 1, 2, 3. \end{aligned} \quad (4.9)$$

Therefore

$$\begin{aligned} p|_{ab} &= b^n{}_{a|nb} \beta + c^n{}_{a|nb} \gamma \\ &\quad + [I_{1,b} b^n{}_{a|n} + (b^n{}_a I_{1,n})|_b] \beta_1 + [I_{2,b} b^n{}_{a|n} + (b^n{}_a I_{2,n})|_b] \beta_2 \\ &\quad + I_{1,n} I_{1,b} b^n{}_a \beta_{11} + I_{2,n} I_{2,b} b^n{}_a \beta_{22} + (I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) b^n{}_a \beta_{12} \\ &\quad + [I_{1,b} c^n{}_{a|n} + (c^n{}_a I_{1,n})|_b] \gamma_1 + [I_{2,b} c^n{}_{a|n} + (c^n{}_a I_{2,n})|_b] \gamma_2 \\ &\quad + I_{1,n} I_{1,b} c^n{}_a \gamma_{11} + I_{2,n} I_{2,b} c^n{}_a \gamma_{22} + (I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) c^n{}_a \gamma_{12} \\ &\quad + [F^{-A}{}_n b^n{}_{a|b} + F^{-A}{}_b b^n{}_{a|n} + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) b^n{}_a] \beta_{,A} \\ &\quad + [F^{-A}{}_n c^n{}_{a|b} + F^{-A}{}_b c^n{}_{a|n} + (F^{-B}{}_b F^{-A}{}_{n,B} - \gamma^m{}_{nb} F^{-A}{}_m) c^n{}_a] \gamma_{,A} \\ &\quad + F^{-A}{}_n b^n{}_a I_{1,b} \beta_{1,A} + F^{-A}{}_n b^n{}_a I_{2,b} \beta_{2,A} + F^{-A}{}_n c^n{}_a I_{1,b} \gamma_{1,A} + F^{-A}{}_n c^n{}_a I_{2,b} \gamma_{2,A} \\ &\quad + F^{-A}{}_n F^{-B}{}_b b^n{}_a \beta_{,AB} + F^{-A}{}_n F^{-B}{}_b c^n{}_a \gamma_{,AB}. \end{aligned} \quad (4.10)$$

The functions β and γ and their derivatives are independent, and hence, the coefficient of each function must

be symmetric. Thus, we have the following set of universality constraints for homogeneous Cauchy elasticity

$$\left\{ \begin{array}{l} \mathcal{A}_{ab}^0 = b_{a|nb}^n, \end{array} \right. \quad (4.11)$$

$$\mathcal{A}_{ab}^1 = I_{1,b} b_{a|n}^n + (b_a^n I_{1,n})_{|b}, \quad (4.12)$$

$$\mathcal{A}_{ab}^2 = I_{2,b} b_{a|n}^n + (b_a^n I_{2,n})_{|b}, \quad (4.13)$$

$$\mathcal{A}_{ab}^{11} = I_{1,n} I_{1,b} b_a^n, \quad (4.14)$$

$$\mathcal{A}_{ab}^{22} = I_{2,n} I_{2,b} b_a^n, \quad (4.15)$$

$$\mathcal{A}_{ab}^{12} = (I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) b_a^n, \quad (4.16)$$

$$\mathcal{B}_{ab}^0 = c_{a|nb}^n, \quad (4.17)$$

$$\mathcal{B}_{ab}^1 = I_{1,b} c_{a|n}^n + (c_a^n I_{1,n})_{|b}, \quad (4.18)$$

$$\mathcal{B}_{ab}^2 = I_{2,b} c_{a|n}^n + (c_a^n I_{2,n})_{|b}, \quad (4.19)$$

$$\mathcal{B}_{ab}^{11} = I_{1,n} I_{1,b} c_a^n, \quad (4.20)$$

$$\mathcal{B}_{ab}^{22} = I_{2,n} I_{2,b} c_a^n, \quad (4.21)$$

$$\mathcal{B}_{ab}^{12} = (I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) c_a^n. \quad (4.22)$$

The six terms highlighted in blue are identical to those of hyperealsticity [Ericksen, 1954]. The following is the set of universal inhomogeneity constraints inhomogeneous Cauchy elasticity

$$\left\{ \begin{array}{l} \mathcal{C}_{ab}^{0A} = F^{-A}_n b_{a|b}^n + F^{-A}_b b_{a|n}^n + (F^{-B}_b F^{-A}_{n,B} - \gamma^m_{nb} F^{-A}_m) b_a^n, \end{array} \right. \quad (4.23)$$

$$\mathcal{C}_{ab}^{1A} = (F^{-A}_n I_{1,b} + F^{-A}_b I_{1,n}) b_a^n, \quad (4.24)$$

$$\mathcal{C}_{ab}^{2A} = (F^{-A}_n I_{2,b} + F^{-A}_b I_{2,n}) b_a^n, \quad (4.25)$$

$$\mathcal{C}_{ab}^{0AB} = (F^{-A}_n F^{-B}_b + F^{-B}_n F^{-A}_b) b_a^n, \quad (4.26)$$

$$\mathcal{D}_{ab}^{0A} = F^{-A}_n c_{a|b}^n + F^{-A}_b c_{a|n}^n + (F^{-B}_b F^{-A}_{n,B} - \gamma^m_{nb} F^{-A}_m) c_a^n, \quad (4.27)$$

$$\mathcal{D}_{ab}^{1A} = (F^{-A}_n I_{1,b} + F^{-A}_b I_{1,n}) c_a^n, \quad (4.28)$$

$$\mathcal{D}_{ab}^{2A} = (F^{-A}_n I_{2,b} + F^{-A}_b I_{2,n}) c_a^n, \quad (4.29)$$

$$\mathcal{D}_{ab}^{0AB} = (F^{-A}_n F^{-B}_b + F^{-B}_n F^{-A}_b) c_a^n. \quad (4.30)$$

These terms must be symmetric in (a, b) for $A = 1, 2, 3$ and $A \leq B = 1, 2, 3$. The six terms highlighted in purple are identical to those of inhomogeneous isotropic hyperelasticity [Yavari, 2021a].

Recall that Ericksen's universality constraints of isotropic hyperelasticity are [Ericksen, 1954]:

$$\left\{ \begin{array}{l} \mathcal{A}_{ab}^1 = b_{a|bn}^n, \end{array} \right. \quad (4.31)$$

$$\mathcal{A}_{ab}^2 = c_{a|bn}^n, \quad (4.32)$$

$$\mathcal{A}_{ab}^{11} = b_{a|n}^n I_{1,b} + (b_a^n I_{1,n})_{|b}, \quad (4.33)$$

$$\mathcal{A}_{ab}^{22} = c_{a|n}^n I_{2,b} + (c_a^n I_{2,n})_{|b}, \quad (4.34)$$

$$\mathcal{A}_{ab}^{12} = (b_a^n I_{2,n})_{|b} + b_{a|n}^n I_{2,b} - [(c_a^n I_{1,n})_{|b} + c_{a|n}^n I_{1,b}], \quad (4.35)$$

$$\mathcal{A}_{ab}^{111} = b_{a|n}^n I_{1,n} I_{1,b}, \quad (4.36)$$

$$\mathcal{A}_{ab}^{222} = c_{a|n}^n I_{2,n} I_{2,b}, \quad (4.37)$$

$$\mathcal{A}_{ab}^{112} = b_a^n (I_{1,b} I_{2,n} + I_{1,n} I_{2,b}) - c_a^n I_{1,n} I_{1,b}, \quad (4.38)$$

$$\mathcal{A}_{ab}^{122} = b_a^n I_{2,b} I_{2,n} - c_a^n (I_{1,b} I_{2,n} + I_{1,n} I_{2,b}). \quad (4.39)$$

Also the universal inhomogeneity constraints of hyperelasticity are [Yavari, 2021a]:

$$\begin{cases} \mathcal{C}_{ab}^{1A} = F^{-A}_n b_a^n|_b + F^{-A}_b b_a^n|_n + b_a^n [F^{-B}_b F^{-A}_{n,B} - \gamma^m_{nb} F^{-A}_m], \end{cases} \quad (4.40)$$

$$\begin{cases} \mathcal{C}_{ab}^{2A} = F^{-A}_n c_a^n|_b + F^{-A}_b c_a^n|_n + c_a^n [F^{-B}_b F^{-A}_{n,B} - \gamma^m_{nb} F^{-A}_m], \end{cases} \quad (4.41)$$

$$\begin{cases} \mathcal{C}_{ab}^{11A} = b_a^n [F^{-A}_n I_{1,b} + F^{-A}_b I_{1,n}], \end{cases} \quad (4.42)$$

$$\begin{cases} \mathcal{C}_{ab}^{22A} = c_a^n [F^{-A}_n I_{2,b} + F^{-A}_b I_{2,n}], \end{cases} \quad (4.43)$$

$$\begin{cases} \mathcal{C}_{ab}^{12A} = b_a^n [F^{-A}_n I_{2,b} + F^{-A}_b I_{2,n}] - c_a^n [F^{-A}_n I_{1,b} + F^{-A}_b I_{1,n}], \end{cases} \quad (4.44)$$

$$\begin{cases} \mathcal{C}_{ab}^{1AB} = b_a^n [F^{-A}_n F^{-B}_b + (F^{-1})^B_n F^{-A}_b], \end{cases} \quad (4.45)$$

$$\begin{cases} \mathcal{C}_{ab}^{2AB} = c_a^n [F^{-A}_n F^{-B}_b + (F^{-1})^B_n F^{-A}_b]. \end{cases} \quad (4.46)$$

First, notice that there are nine sets of universality constraints (4.31)-(4.39) in homogeneous isotropic hyperelasticity [Ericksen, 1954] compared to twelve in compressible isotropic Cauchy elasticity (4.11)-(4.22). Second, the six terms shown in blue are identical to those of hyperelasticity. Thus, we first look at these six terms.

Ericksen [1954] used the following result. Suppose \mathbf{u} and \mathbf{v} are vectors such that $\mathbf{u} \otimes \mathbf{v} = \mathbf{v} \otimes \mathbf{u}$. Assuming that $\mathbf{v} \neq \mathbf{0}$ (if both vectors are zero this equality trivially holds) one can write

$$\mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{v}|^2} \mathbf{v} = \lambda \mathbf{v}, \quad (4.47)$$

i.e., \mathbf{u} and \mathbf{v} must be parallel. Now suppose that $\mathbf{u}^\flat = d\phi$ and $\mathbf{v}^\flat = d\psi$, where \flat is the flat operator that gives the 1-form corresponding to a vector, d is the exterior derivative, and ϕ and ψ are scalar fields. Note that $\lambda = \lambda(\mathbf{u}, \mathbf{v}) = \lambda(\phi, \psi)$. Thus, $\mathbf{u}^\flat = d\phi = \lambda(\phi, \psi)d\psi$. Hence

$$0 = d \circ d\phi = d\lambda(\phi, \psi) \wedge d\psi = \frac{\partial \lambda}{\partial \phi} d\phi \wedge d\psi, \quad (4.48)$$

where \wedge is the wedge product of differential forms. Therefore, $\lambda = \lambda(\psi)$, i.e., $\mathbf{u}^\flat = \lambda(\psi)d\psi$.

W_{111} (\mathcal{A}_{ab}^{11}) and W_{222} (\mathcal{B}_{ab}^{22}) terms: Symmetry of the coefficient of the \mathcal{A}_{ab}^{11} term implies that [Ericksen, 1954]

$$(b_a^n I_{1,n}) I_{1,b} = (b_b^n I_{1,n}) I_{1,a}. \quad (4.49)$$

One concludes that either $I_{1,a} = 0$ or $b_a^n I_{1,n} = A^{-1} I_{1,a}$, for some scalar function A . This means that $I_{1,a}$ is an eigenvector of b_a^n with eigenvalue A^{-1} . We know that $b^{am} c_{mb} = \delta_b^a$, and hence, $b_b^m c_m^a = \delta_b^a$. Similarly, $b_{am} c^{mb} = \delta_a^b$, and hence, $b_m^a c_b^m = \delta_b^a$. Therefore, $c_a^n I_{1,n} = A I_{1,a}$.⁹ Similarly, the symmetry of the coefficient of the \mathcal{B}_{ab}^{22} term implies that

$$(c_a^n I_{2,n}) I_{2,b} = (c_b^n I_{2,n}) I_{2,a}. \quad (4.50)$$

The conclusion is that either $I_{2,a} = 0$ or $c_a^n I_{2,n} = \bar{A} I_{2,a}$, for some scalar function \bar{A} . One also has $b_a^n I_{2,n} = \bar{A}^{-1} I_{2,a}$. In summary

$$\begin{cases} c_a^n I_{1,n} = A I_{1,a}, \end{cases} \quad (4.51)$$

$$\begin{cases} c_a^n I_{2,n} = \bar{A} I_{2,a}, \end{cases} \quad (4.52)$$

$$\begin{cases} b_a^n I_{1,n} = A^{-1} I_{1,a}, \end{cases} \quad (4.53)$$

$$\begin{cases} b_a^n I_{2,n} = \bar{A}^{-1} I_{2,a}. \end{cases} \quad (4.54)$$

⁹Note that both \mathbf{b} and \mathbf{c} are positive-definite. This means that when I_1 is not constant, $A \neq 0$, and hence, A^{-1} is an eigenvalue of \mathbf{c} .

\mathcal{A}_{ab}^{22} and \mathcal{B}_{ab}^{11} terms: Symmetry of the coefficient of the \mathcal{A}_{ab}^{22} term implies that

$$(b_a^n I_{2,n}) I_{2,b} = (b_b^n I_{2,n}) I_{2,a}, \quad \text{or} \quad \bar{\mathbf{A}}^{-1} I_{2,a} I_{2,b} = \bar{\mathbf{A}}^{-1} I_{2,b} I_{2,a}, \quad (4.55)$$

which is trivially satisfied. Similarly, the symmetry of the coefficient of the \mathcal{B}_{ab}^{11} term implies that

$$(c_a^n I_{1,n}) I_{1,b} = (c_b^n I_{1,n}) I_{1,a}, \quad \text{or} \quad \mathbf{A} I_{1,a} I_{1,b} = \mathbf{A} I_{1,b} I_{1,a}, \quad (4.56)$$

which is also trivially satisfied.

\mathcal{A}_{ab}^{12} and \mathcal{B}_{ab}^{12} terms: Symmetry of the coefficient of the \mathcal{A}_{ab}^{12} term implies that

$$(I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) b_a^n = (I_{1,n} I_{2,a} + I_{2,n} I_{1,a}) b_b^n, \quad (4.57)$$

which is identical to the symmetry condition that comes from the coefficient of the W_{112} in hyperelasticity [Ericksen, 1954]. Thus

$$(I_{1,b} I_{2,n} + I_{1,n} I_{2,b}) b_a^n = (I_{1,a} I_{2,n} + I_{1,n} I_{2,a}) b_b^n. \quad (4.58)$$

Hence

$$I_{1,b} \bar{\mathbf{A}}^{-1} I_{2,a} + \mathbf{A}^{-1} I_{1,a} I_{2,b} = I_{1,a} \bar{\mathbf{A}}^{-1} I_{2,b} + \mathbf{A}^{-1} I_{1,b} I_{2,a}. \quad (4.59)$$

Or

$$(\mathbf{A}^{-1} - \bar{\mathbf{A}}^{-1}) (I_{1,a} I_{2,b} - I_{1,b} I_{2,a}) = 0. \quad (4.60)$$

Similarly, the symmetry of the coefficient of the \mathcal{B}_{ab}^{12} term implies that

$$(I_{1,n} I_{2,b} + I_{2,n} I_{1,b}) c_a^n = (I_{1,n} I_{2,a} + I_{2,n} I_{1,a}) c_b^n. \quad (4.61)$$

This is simplified to read

$$(\mathbf{A} - \bar{\mathbf{A}}) (I_{1,a} I_{2,b} - I_{1,b} I_{2,a}) = 0, \quad (4.62)$$

which is identical to the symmetry condition that comes from the coefficient of the W_{221} in hyperelasticity [Ericksen, 1954]. If either I_1 or I_2 is constant, (4.60) is satisfied for any \mathbf{A} and $\bar{\mathbf{A}}$. If $\bar{\mathbf{A}} \neq \mathbf{A}$, one has $I_{1,a} I_{2,b} = I_{1,b} I_{2,a}$, which implies that $I_{1,a}$ and $I_{2,a}$ are parallel. However, these are eigenvectors of b_a^b corresponding to the distinct eigenvalues $\bar{\mathbf{A}} \neq \mathbf{A}$ and must be perpendicular. This contradiction shows that $\bar{\mathbf{A}} = \mathbf{A}$.

Next, Ericksen shows that I_1 and I_2 must be functionally dependent.¹⁰ If either I_1 or I_2 is constant, this is trivially the case. If all the eigenvalues of \mathbf{b} (and hence those of \mathbf{c}) are distinct, each must have a one-dimensional eigenspace. Therefore, $I_{1,a}$ and $I_{2,a}$, which are eigenvectors corresponding to the same eigenvalue, must be parallel. This guarantees that the rank of the Jacobian matrix in (4.63) is less than 2, and hence, I_1 and I_2 are functionally dependent. Note that

$$I_1 = \frac{1}{c_1} + \frac{1}{c_2} + \frac{1}{c_3}, \quad I_2 = \frac{1}{c_1 c_2} + \frac{1}{c_2 c_3} + \frac{1}{c_3 c_1}, \quad c_1 c_2 c_3 = 1, \quad (4.64)$$

where c_i are eigenvalues of \mathbf{c} . If two eigenvalues are equal, e.g., $c_1 = c_2$, one has (the case $c_1 = c_2 = c_3$ is trivial as incompressibility implies $c_1 = c_2 = c_3 = 1$.)

$$I_1 = \frac{2}{c_1} + c_1^2, \quad I_2 = \frac{1}{c_1^2} + 2c_1. \quad (4.65)$$

Clearly, in this case I_1 and I_2 are functionally dependent. Therefore, there is a scalar function \mathbf{B} such that, $I_1 = I_1(\mathbf{B})$ and $I_2 = I_2(\mathbf{B})$. Similarly, $c_i = c_i(\mathbf{B})$.

¹⁰ I_1 and I_2 are functionally dependent if there exists a non-trivial function (a function that is not identically zero) such that $F(I_1, I_2) = 0$. Taking derivatives one obtains

$$\begin{bmatrix} I_{1,1} & I_{2,1} \\ I_{1,2} & I_{2,2} \\ I_{1,3} & I_{2,3} \end{bmatrix} \begin{bmatrix} \frac{\partial F}{\partial I_1} \\ \frac{\partial F}{\partial I_2} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}. \quad (4.63)$$

For F to be non-trivial the Jacobian matrix must have rank less than 2.

W_{11} (\mathcal{A}_{ab}^1) and W_{22} (\mathcal{B}_{ab}^2) terms: Note that $I_{i,a} = I'_i(\mathbf{B})\mathbf{B}_{,a}$, $i = 1, 2$. The second term in the coefficient of W_{22} (\mathcal{B}_{ab}^2) is simplified to read $(c_a^n I_{2,n})_{,b} = (c_1 I'_2 \mathbf{B}_{,a})_{,b} = (c_1 I'_2)' \mathbf{B}_{,b} \mathbf{B}_{,a} + c_1 I'_2 \mathbf{B}_{,ab}$, which is symmetric. Therefore, symmetry of the coefficient of W_{22} implies that

$$I'_2(c_{a|n}^n \mathbf{B}_{,b} - c_{b|n}^n \mathbf{B}_{,a}) = 0. \quad (4.66)$$

Thus, either I_2 is constant or

$$c_{a,n}^n = D \mathbf{B}_{,a}, \quad (4.67)$$

where D is a scalar field. Similarly, symmetry of the coefficient of W_{11} (\mathcal{A}_{ab}^1) dictates that either I_1 is constant or

$$b_{a|n}^n = E \mathbf{B}_{,a}, \quad (4.68)$$

where E is a scalar field.

\mathcal{A}_{ab}^2 and \mathcal{B}_{ab}^1 terms: The symmetry of \mathcal{A}_{ab}^2 terms implies that

$$I'_2(b_{a|n}^n \mathbf{B}_{,b} - b_{b|n}^n \mathbf{B}_{,a}) = I'_2(E \mathbf{B}_{,a} \mathbf{B}_{,b} - E \mathbf{B}_{,b} \mathbf{B}_{,a}) = 0, \quad (4.69)$$

which is trivially satisfied. Similarly, the symmetry of \mathcal{B}_{ab}^1 terms implies that

$$I'_1(c_{a|n}^n \mathbf{B}_{,b} - c_{b|n}^n \mathbf{B}_{,a}) = I'_1(D \mathbf{B}_{,a} \mathbf{B}_{,b} - D \mathbf{B}_{,b} \mathbf{B}_{,a}) = 0, \quad (4.70)$$

which is also trivially satisfied.

The only two remaining universality constraints are \mathcal{A}_{ab}^0 and \mathcal{B}_{ab}^0 terms, which are identical to the coefficients of W_1 and W_2 in hyperelasticity. Using this two sets of universality constraints, [Ericksen \[1954\]](#) found Families 1-4 of universal deformations.

Note that if I_1 and I_2 are constant, (4.11)-(4.22) reduce to the symmetry of the two terms $b_{a|nb}^n$ and $c_{a|nb}^n$, which is exactly what one would have in hyperelasticity. We have thus shown that although there are more universality constraints in incompressible isotropic Cauchy elasticity and some of the universality constraints are not identical to those of hyperelasticity, nevertheless the two sets of universality constraints are equivalent.

Next we show that the universal inhomogeneity constraints of Cauchy elasticity (4.23)-(4.30) are equivalent to those of hyperelasticity (4.40)-(4.46). We only need to check the terms in black, as the ones highlighted in purple are identical. Let us define $\mathbf{P}_{bn}^A = F^{-A}_n I_{1,b} + F^{-A}_b I_{1,n}$ and note that $\mathbf{P}_{nb}^A = \mathbf{P}_{bn}^A$. From (4.24) we know that

$$\mathbf{P}_{bn}^A b_a^n = \mathbf{P}_{an}^A b_b^n. \quad (4.71)$$

Multiply both sides by $c_m^a c_k^b$ to obtain $c_k^b \mathbf{P}_{bn}^A b_a^n c_m^a = c_m^a \mathbf{P}_{an}^A b_b^n c_k^b$, or $c_k^b \mathbf{P}_{bn}^A \delta_m^n = c_m^a \mathbf{P}_{an}^A \delta_k^n$. Thus

$$c_k^b \mathbf{P}_{bm}^A = c_m^a \mathbf{P}_{ak}^A, \quad \text{or} \quad c_a^n \mathbf{P}_{nb}^A = c_b^n \mathbf{P}_{na}^A, \quad (4.72)$$

which is (4.28). Similarly, (4.29) is equivalent to (4.25). Moreover, symmetry of the terms (4.42) and (4.43) imply the symmetry of (4.44). Therefore, we have proved the following result.

Proposition 4.1. *The set of universal deformations of incompressible isotropic Cauchy elasticity is identical to that of incompressible isotropic hyperelasticity. For a given universal deformation, the corresponding set of universal inhomogeneities is identical to that of hyperelasticity.*

5 Conclusions

The existing studies of universal deformations have been restricted to hyperelasticity, or more generally, to materials that have an underlying energy function. In this paper, we extended the analysis of universal deformations to Cauchy elastic solids, which in general, do not have an energy function. We considered both compressible and incompressible inhomogeneous isotropic Cauchy elasticity. As hyperelasticity is a

proper subset of Cauchy elasticity, one would expect the set of universal deformations of Cauchy elasticity to be a proper subset of that of hyperelasticity. We proved the somewhat unexpected result that the sets of universal deformations of isotropic Cauchy elasticity and isotropic hyperelasticity are identical, in both the compressible and incompressible cases. We also proved that their corresponding universal inhomogeneities are identical as well.

Acknowledgments

This work was supported by NSF – Grant No. CMMI 1939901.

References

- J. E. Adkins and R. S. Rivlin. Large elastic deformations of isotropic materials X. Reinforcement by inextensible cords. *Philosophical Transactions of the Royal Society of London. Series A*, 248(944):201–223, 1955.
- M. F. Beatty. General solutions in the equilibrium theory of inextensible elastic materials. *Acta Mechanica*, 29(1):119–126, 1978.
- D. E. Beskos. Universal solutions for fiber-reinforced compressible isotropic elastic materials. *Journal of Elasticity*, 2(3):153–168, 1972.
- J.-P. Boehler. On irreducible representations for isotropic scalar functions. *Zeitschrift für Angewandte Mathematik und Mechanik*, 57(6):323–327, 1977.
- G. Bordiga, A. Piccolroaz, and D. Bigoni. A way to hypo-elastic artificial materials without a strain potential and displaying flutter instability. *Journal of the Mechanics and Physics of Solids*, 158:104665, 2022.
- M. M. Carroll. Controllable deformations of incompressible simple materials. *International Journal of Engineering Science*, 5(6):515–525, 1967.
- M. M. Carroll. Controllable states of stress for compressible elastic solids. *Journal of Elasticity*, 3:57–61, 1973.
- M. M. Carroll. Must elastic materials be hyperelastic? *Mathematics and Mechanics of Solids*, 14(4):369–376, 2009.
- J. Casey. A remark on Cauchy-elasticity. *International Journal of Non-Linear Mechanics*, 40(2-3):331–339, 2005.
- A.-L. Cauchy. Sur les équations qui expriment les conditions d’équilibre ou les lois du mouvement intérieur d’un corps solide, élastique ou non élastique. *Exercices de Mathématiques*, 3:160–187, 1828.
- H. Chi, C. Talischi, O. Lopez-Pamies, and G. H. Paulino. Polygonal finite elements for finite elasticity. *International Journal for Numerical Methods in Engineering*, 101(4):305–328, 2015.
- R. De Pascalis, K. Rajagopal, and G. Saccomandi. Remarks on the use and misuse of the semi-inverse method in the nonlinear theory of elasticity. *Quarterly journal of mechanics and applied mathematics*, 62(4):451–464, 2009.
- T. Doyle and J. L. Ericksen. Nonlinear elasticity. *Advances in Applied Mechanics*, 4:53–115, 1956.
- E. Dragoni. The radial compaction of a hyperelastic tube as a benchmark in compressible finite elasticity. *International Journal of Non-Linear Mechanics*, 31(4):483–493, 1996.
- C. Eckart. The thermodynamics of irreversible processes. 4. The theory of elasticity and anelasticity. *Physical Review*, 73(4):373–382, 1948.

- J. L. Ericksen. Deformations possible in every isotropic, incompressible, perfectly elastic body. *Zeitschrift für Angewandte Mathematik und Physik*, 5(6):466–489, 1954.
- J. L. Ericksen. Deformations possible in every compressible, isotropic, perfectly elastic material. *Studies in Applied Mathematics*, 34(1-4):126–128, 1955.
- J. L. Ericksen. Stress deformation relations for solids. *Canadian Journal of Physics*, 34(2):226–227, 1956.
- J. L. Ericksen and R. S. Rivlin. Large elastic deformations of homogeneous anisotropic materials. *Journal of Rational Mechanics and Analysis*, 3:281–301, 1954.
- R. L. Fosdick. Remarks on Compatibility. *Modern Developments in the Mechanics of Continua*, pages 109–127, 1966.
- R. L. Fosdick. Statically possible radially symmetric deformations in isotropic, incompressible elastic solids. *Zeitschrift für angewandte Mathematik und Physik*, 22:590–607, 1971.
- R. L. Fosdick and K. W. Schuler. On Ericksen’s problem for plane deformations with uniform transverse stretch. *International Journal of Engineering Science*, 7(2):217–233, 1969.
- M. Fruchart, C. Scheibner, and V. Vitelli. Odd viscosity and odd elasticity. *Annual Review of Condensed Matter Physics*, 14:471–510, 2023.
- B. Gairola. *Nonlinear elastic problems*. F.R.N. Nabarro (Ed.), Dislocations in Solids. North-Holland Publishing Co., Amsterdam, 1979.
- A. Golgoon and A. Yavari. Nonlinear elastic inclusions in anisotropic solids. *Journal of Elasticity*, 130(2):239–269, 2018a.
- A. Golgoon and A. Yavari. Line and point defects in nonlinear anisotropic solids. *Zeitschrift für angewandte Mathematik und Physik*, 69:1–28, 2018b.
- A. Golgoon and A. Yavari. On Hashin’s hollow cylinder and sphere assemblages in anisotropic nonlinear elasticity. *Journal of Elasticity*, 146(1):65–82, 2021.
- C. Goodbrake, A. Yavari, and A. Goriely. The anelastic Ericksen problem: Universal deformations and universal eigenstrains in incompressible nonlinear anelasticity. *Journal of Elasticity*, 142(2):291–381, 2020.
- A. Goriely. *The Mathematics and Mechanics of Biological Growth*, volume 45. Springer, 2017.
- A. E. Green and P. M. Naghdi. On thermodynamics, rate of work and energy. *Archive for Rational Mechanics and Analysis*, 40:37–49, 1971.
- M. E. Gurtin. The linear theory of elasticity. In *Handbuch der Physik, Band VIa/2*. Springer-Verlag, Berlin, 1972.
- Z. Hashin. Large isotropic elastic deformation of composites and porous media. *International Journal of Solids and Structures*, 21(7):711–720, 1985.
- C. B. Kafadar. On Ericksen’s problem. *Archive for Rational Mechanics and Analysis*, 47:15–27, 1972.
- W. W. Klingbeil and R. T. Shield. On a class of solutions in plane finite elasticity. *Zeitschrift für angewandte Mathematik und Physik*, 17(4):489–511, 1966.
- J. K. Knowles. Universal states of finite anti-plane shear: Ericksen’s problem in miniature. *The American Mathematical Monthly*, 86(2):109–113, 1979.
- M. Kružík and T. Roubíček. *Mathematical Methods in Continuum Mechanics of Solids*. Springer, 2019.

- A. Kumar and A. Yavari. Nonlinear mechanics of remodeling. *Journal of the Mechanics and Physics of Solids*, 181:105449, 2023.
- V. Lee and K. Bhattacharya. Universal deformations of incompressible nonlinear elasticity as applied to ideal liquid crystal elastomers. *Journal of Elasticity*, pages 1–27, 2023.
- O. Lopez-Pamies, J. Moraleda, J. Segurado, and J. Llorca. On the extremal properties of Hashin’s hollow cylinder assemblage in nonlinear elasticity. *Journal of Elasticity*, 107:1–10, 2012.
- A. Marris. Universal deformations in incompressible isotropic elastic materials. *Journal of Elasticity*, 5(2): 111–128, 1975.
- A. Marris. Two new theorems on Ericksen’s problem. *Archive for Rational Mechanics and Analysis*, 79: 131–173, 1982.
- A. Marris and J. Shiau. Universal deformations in isotropic incompressible hyperelastic materials when the deformation tensor has equal proper values. *Archive for Rational Mechanics and Analysis*, 36(2):135–160, 1970.
- J. E. Marsden and T. J. R. Hughes. *Mathematical Foundations of Elasticity*. Dover Publications, New York, 1994.
- L. A. Mihai and A. Goriely. Controllable deformations of unconstrained ideal nematic elastomers. *Journal of Elasticity*, pages 1–12, 2023.
- A. J. A. Morgan. Some properties of media defined by constitutive equations in implicit form. *International Journal of Engineering Science*, 4(2):155–178, 1966.
- W. Noll. A mathematical theory of the mechanical behavior of continuous media. *Archive for Rational Mechanics and Analysis*, 2(1):197–226, 1958.
- R. W. Ogden. *Non-Linear Elastic Deformations*. Dover, 1984.
- A. C. Pipkin. Generalized plane deformations of ideal fiber-reinforced materials. *Quarterly of Applied Mathematics*, 32(3):253–263, 1974.
- A. C. Pipkin. Stress analysis for fiber-reinforced materials. In *Advances in Applied Mechanics*, volume 19, pages 1–51. Elsevier, 1979.
- A. C. Pipkin and T. G. Rogers. Plane deformations of incompressible fiber-reinforced materials. *Journal of Applied Mechanics*, 38(3):634–640, 1971.
- D. A. Polignone and C. O. Horgan. Pure torsion of compressible non-linearly elastic circular cylinders. *Quarterly of applied mathematics*, 49(3):591–607, 1991.
- S. P. Pradhan and A. Yavari. Accretion–ablation mechanics. *Philosophical Transactions of the Royal Society A*, 381(2263):20220373, 2023.
- K. R. Rajagopal. On implicit constitutive theories. *Applications of Mathematics*, 48:279–319, 2003.
- K. R. Rajagopal. The elasticity of elasticity. *Zeitschrift für angewandte Mathematik und Physik*, 58:309–317, 2007.
- K. R. Rajagopal. Conspectus of concepts of elasticity. *Mathematics and Mechanics of Solids*, 16(5):536–562, 2011.
- R. S. Rivlin. Large elastic deformations of isotropic materials IV. Further developments of the general theory. *Philosophical Transactions of the Royal Society of London A*, 241(835):379–397, 1948.

- R. S. Rivlin. Large elastic deformations of isotropic materials. V. The problem of flexure. *Proceedings of the Royal Society of London A*, 195(1043):463–473, 1949a.
- R. S. Rivlin. A note on the torsion of an incompressible highly elastic cylinder. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 45, pages 485–487. Cambridge University Press, 1949b.
- R. S. Rivlin. Reflections on certain aspects of themomechanics. In *Contributi del Centro Linceo Interdisciplinare di Scienze Matematiche e loro Applicazioni*, pages 11–43. 1986.
- R. S. Rivlin and J. L. Ericksen. Stress-deformation relations for isotropic materials. *Journal of Rational Mechanics and Analysis*, 4:323–425, 1955.
- R. S. Rivlin and D. W. Saunders. Large elastic deformations of isotropic materials VII. Experiments on the deformation of rubber. *Philosophical Transactions of the Royal Society of London A*, 243(865):251–288, 1951.
- G. Saccomandi. Universal solutions and relations in finite elasticity. In *Topics in finite elasticity*, pages 95–130. Springer, 2001.
- S. Sadik and A. Yavari. Nonlinear anisotropic viscoelasticity. *Journal of the Mechanics and Physics of Solids*, 182:105461, 2024.
- C. Scheibner, A. Souslov, D. Banerjee, P. Surówka, W. T. Irvine, and V. Vitelli. Odd elasticity. *Nature Physics*, 16(4):475–480, 2020.
- M. F. Shojaei and A. Yavari. Compatible-strain mixed finite element methods for incompressible nonlinear elasticity. *Journal of Computational Physics*, 361:247–279, 2018.
- M. Singh and A. C. Pipkin. Note on Ericksen’s problem. *Zeitschrift für angewandte Mathematik und Physik*, 16(5):706–709, 1965.
- E. B. Tadmor, R. E. Miller, and R. S. Elliott. *Continuum Mechanics and Thermodynamics: From Fundamental Concepts to Governing Equations*. Cambridge University Press, 2012.
- C. Truesdell. The mechanical foundations of elasticity and fluid dynamics. *Journal of Rational Mechanics and Analysis*, 1(1):125–300, 1952.
- C. Truesdell. *The Elements of Continuum Mechanics*. Springer-Verlag, 1966.
- C. Truesdell and W. Noll. *The Non-Linear Field Theories of Mechanics*. Springer, 2004.
- C. C. Wang. On representations for isotropic functions: Part I. Isotropic functions of symmetric tensors and vectors. *Archive for Rational Mechanics and Analysis*, 33:249–267, 1969.
- Z. Wesolowski and A. Seeger. On the screw dislocation in finite elasticity. In *Mechanics of Generalized Continua. Proceedings of the IUTAM Symposium on the Generalized Cosserat Continuum and the Continuum Theory of Dislocations with Applications (Ed. E. Kröner, Springer, Berlin, etc., 1968)*, pages 294–297. Springer, 1968.
- A. Yavari. Universal deformations in inhomogeneous isotropic nonlinear elastic solids. *Proceedings of the Royal Society A*, 477(2253):20210547, 2021a.
- A. Yavari. On Eshelby’s inclusion problem in nonlinear anisotropic elasticity. *Journal of Micromechanics and Molecular Physics*, 6(01):2150002, 2021b.
- A. Yavari. Universal displacements in inextensible fiber-reinforced linear elastic solids. *Mathematics and Mechanics of Solids*, pages 1–17, 2023. doi: 10.1177/10812865231181924.
- A. Yavari and A. Goriely. Riemann–Cartan geometry of nonlinear dislocation mechanics. *Archive for Rational Mechanics and Analysis*, 205(1):59–118, 2012a.

- A. Yavari and A. Goriely. Weyl geometry and the nonlinear mechanics of distributed point defects. *Proceedings of the Royal Society A*, 468(2148):3902–3922, 2012b.
- A. Yavari and A. Goriely. Riemann–Cartan geometry of nonlinear disclination mechanics. *Mathematics and Mechanics of Solids*, 18(1):91–102, 2013a.
- A. Yavari and A. Goriely. Nonlinear elastic inclusions in isotropic solids. *Proc. R. Soc. A*, 469(2160):20130415, 2013b.
- A. Yavari and A. Goriely. The geometry of discombinations and its applications to semi-inverse problems in anelasticity. *Proceedings of the Royal Society A*, 470(2169):20140403, 2014.
- A. Yavari and A. Goriely. The twist-fit problem: Finite torsional and shear eigenstrains in nonlinear elastic solids. *Proceedings of the Royal Society A*, 471(2183):20150596, 2015.
- A. Yavari and A. Goriely. The anelastic Ericksen problem: Universal eigenstrains and deformations in compressible isotropic elastic solids. *Proc. R. Soc. A*, 472(2196):20160690, 2016.
- A. Yavari and A. Goriely. Universal deformations in anisotropic nonlinear elastic solids. *Journal of the Mechanics and Physics of Solids*, 156:104598, 2021.
- A. Yavari and A. Goriely. Universality in anisotropic linear anelasticity. *Journal of Elasticity*, 150(2):241–259, 2022a.
- A. Yavari and A. Goriely. The universal program of linear elasticity. *Mathematics and Mechanics of Solids*, 2022b.
- A. Yavari and A. Goriely. The universal program of nonlinear hyperelasticity. *Journal of Elasticity*, 154(1):91–146, 2023.
- A. Yavari and S. P. Pradhan. Accretion mechanics of nonlinear elastic circular cylindrical bars under finite torsion. *Journal of Elasticity*, 152(1-2):29–60, 2022.
- A. Yavari, C. Goodbrake, and A. Goriely. Universal displacements in linear elasticity. *Journal of the Mechanics and Physics of Solids*, 135:103782, 2020.
- A. Yavari, Y. Safa, and A. Soleiman Fallah. Finite extension of accreting nonlinear elastic solid circular cylinders. *Continuum Mechanics and Thermodynamics*, pages 1–17, 2023.
- L. M. Zubov. *Nonlinear Theory of Dislocations and Disclinations in Elastic Bodies*, volume 47. Springer Science & Business Media, 1997.