

Controllable Deformations in Compressible Isotropic Implicit Elasticity*

Arash Yavari^{†1,2} and Alain Goriely³

¹*School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

²*The George W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA*

³*Mathematical Institute, University of Oxford, Oxford, OX2 6GG, UK*

July 24, 2024

Abstract

For a given material, *controllable deformations* are those deformations that can be maintained in the absence of body forces and by applying only boundary tractions. For a given class of materials, *universal deformations* are those deformations that are controllable for any material within the class. In this paper, we characterize the universal deformations in compressible isotropic implicit elasticity defined by solids whose constitutive equations, in terms of the Cauchy stress $\boldsymbol{\sigma}$ and the left Cauchy-Green strain \mathbf{b} , have the implicit form $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$. We prove that universal deformations are homogeneous. However, an important observation is that, unlike Cauchy (and Green) elasticity, not every homogeneous deformation is constitutively admissible for a given implicit-elastic solid. In other words, the set of universal deformations is material-dependent, yet it remains a subset of homogeneous deformations.

Keywords: Universal deformation, implicit constitutive equations, Cauchy elasticity, hyperelasticity, Green elasticity, implicit elasticity, isotropic solids.

1 Introduction

For a specific class of materials, universal deformations are those deformations that can be maintained in the absence of body forces, solely by applying boundary tractions, for all members of the material class. For Cauchy elastic (and particularly, hyperelastic) solids, universal deformations are independent of the specific material within the class. However, the boundary tractions required to maintain a universal deformation explicitly depend on the particular material. Universal deformations have played a pivotal role in nonlinear elasticity and anelasticity. The following are some examples:

- They have played a crucial organizational role in the semi-inverse solutions in nonlinear elasticity [Knowles, 1979, Polignone and Horgan, 1991, De Pascalis et al., 2009, Tadmor et al., 2012, Goriely, 2017], and more recently in anelasticity [Kumar and Yavari, 2023] and viscoelasticity [Sadik and Yavari, 2024].
- They provide guidance for designing experiments aimed at determining the constitutive relations of a specific material [Rivlin and Saunders, 1951, Doyle and Ericksen, 1956, Saccomandi, 2001a].
- All the existing exact solutions for defects in nonlinear solids are associated with universal deformations [Wesolowski and Seeger, 1968, Gairola, 1979, Zubov, 1997, Yavari and Goriely, 2012a,b, 2013a, 2014, Golgoon and Yavari, 2018b].

*To appear in *Zeitschrift für Angewandte Mathematik und Physik*.

[†]Corresponding author, e-mail: arash.yavari@ce.gatech.edu

- Universal deformations have been crucial in finding exact solutions for the stress fields of distributed finite eigenstrains in nonlinear solids, as well as in solving the nonlinear analogues of Eshelby’s inclusion problem [Yavari and Goriely, 2013b, 2015, Golgoon and Yavari, 2018a, Yavari, 2021b].
- Universal deformations are exact solutions that have served as benchmark problems in computational mechanics [Dragoni, 1996, Saccomandi, 2001a, Chi et al., 2015, Shojaei and Yavari, 2018].
- Universal deformations have been utilized in deriving effective properties for nonlinear composites [Hashin, 1985, Lopez-Pamies et al., 2012, Golgoon and Yavari, 2021].

The systematic investigation of universal deformations for homogeneous compressible and incompressible isotropic hyperelastic solids began in the 1950s with the seminal works of Jerry Ericksen [Ericksen, 1954, 1955] whose work was inspired by the earlier contributions of Ronald Rivlin [Rivlin, 1948, 1949a,b]. Ericksen [1955] demonstrated that in homogeneous compressible isotropic solids, universal deformations are homogeneous. Characterizing universal deformations in the presence of internal constraints such as incompressibility poses a particularly challenging problem [Saccomandi, 2001a]. Ericksen [1954] identified four families of universal deformations for incompressible isotropic hyperelastic solids. He conjectured that a deformation with constant principal invariants is homogeneous, but this conjecture turned out to be incorrect [Fosdick, 1966]. Later on, a fifth family of universal deformations was found [Singh and Pipkin, 1965, Klingbeil and Shield, 1966]. The fifth family of universal deformations are inhomogeneous, yet these deformations have constant principal invariants. To this date, the problem of whether there exist additional inhomogeneous constant-principal invariant universal deformations remains open (Ericksen’s problem).

The investigation into universal deformations was expanded to inhomogeneous anisotropic solids in [Yavari and Goriely, 2021, Yavari, 2021a, Yavari and Goriely, 2023]. Before our studies, there had been limited research on universal deformations in anisotropic solids [Ericksen and Rivlin, 1954]. The comprehensive analyses presented in [Yavari and Goriely, 2021, Yavari, 2021a, Yavari and Goriely, 2023] encompassed both compressible and incompressible isotropic, transversely isotropic, orthotropic, and monoclinic solids. It was demonstrated that for these three classes of compressible anisotropic solids, universal deformations are homogeneous, and the preferred material directions are uniform. Moreover, for isotropic solids and each of the three classes of anisotropic solids, the corresponding universal inhomogeneities, which represent inhomogeneities in the energy function compatible with the universality constraints, were characterized. For inhomogeneous incompressible isotropic and the three classes of inhomogeneous incompressible anisotropic solids, the corresponding universal inhomogeneities for each of the above six known families of universal deformations were determined.

In linear elasticity, universal displacements are counterparts to universal deformations [Truesdell, 1966, Gurtin, 1972, Carroll, 1973, Yavari et al., 2020]. Yavari et al. [2020] showed the explicit dependence of universal displacements on the symmetry class of the material. In particular, the larger the symmetry group, the larger the corresponding set of universal displacements. The study of universal displacements has also been expanded to include inhomogeneous solids [Yavari and Goriely, 2022b], linear anelasticity [Yavari and Goriely, 2022a], and compressible anisotropic linear elastic solids reinforced by a single family of inextensible fibers [Yavari, 2023].

Recently, there have been extensions of Ericksen’s analysis to include anelasticity. Yavari and Goriely [2016] proved that in compressible anelasticity, universal deformations are covariantly homogeneous, and for simply-connected bodies universal eigenstrains are impotent (zero-stress). A partial characterization of universal deformations and eigenstrains in incompressible anelasticity was given in [Goodbrake et al., 2020]. In particular, it was observed that the six known families of universal deformations for incompressible, isotropic elastic solids are invariant under specific Lie subgroups of the special Euclidean group. There are also recent studies on universal deformations and eigenstrains in accreting bodies [Yavari and Pradhan, 2022, Yavari et al., 2023, Pradhan and Yavari, 2023] and investigations into universal deformations in liquid crystal elastomers [Lee and Bhattacharya, 2023, Mihai and Goriely, 2023].

In a recent study, Yavari [2024] extended the analysis of universal deformations and inhomogeneities to inhomogeneous compressible and incompressible isotropic Cauchy elasticity. In Cauchy elasticity, which includes hyperelasticity (Green elasticity) as a special case, an energy function does not generally exist. It was demonstrated that somewhat unexpectedly the sets of universal deformations and inhomogeneities of Cauchy elasticity are identical to those of Green elasticity in both compressible and incompressible cases.

Here, we study universal deformations in compressible isotropic solids with implicit constitutive equations of the form $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$, where $\boldsymbol{\sigma}$ is the Cauchy stress and \mathbf{b} is the left Cauchy-Green strain. This class of materials was introduced by Morgan [1966] who used a result due to Rivlin and Ericksen [1955] to simplify such implicit constitutive equations for isotropic solids. This work remained largely unnoticed until Rajagopal and his collaborators started a major research program on the mechanics of elastic solids with implicit constitutive equations in the past two decades [Rajagopal, 2003, 2007, Bustamante, 2009, Bustamante and Rajagopal, 2011]. There have been recent studies exploring universal displacements and deformations for subclasses of incompressible implicit isotropic elasticity. Bustamante [2020b] considered incompressible isotropic elastic solids for which the linearized strain is a function of Cauchy stress. He showed that inflation and uniform extension/compression of a cylindrical annulus is a universal displacement for this class of implicit elastic solids. Bustamante [2020a] showed that the known universal deformations of incompressible isotropic hyperelasticity are also universal for the subclass of incompressible isotropic implicit elastic solids for which the Hencky strain is a function of the Kirchhoff stress.

This paper is organized as follows. In §2, implicit elasticity is briefly reviewed. Universal deformations of compressible isotropic implicit-elastic solids are characterized in §3. Conclusions are given in §4.

2 Implicit elasticity

Consider a body that in its undeformed configuration is identified with an embedded submanifold \mathcal{B} of the Euclidean ambient space \mathcal{S} . The flat metric of the Euclidean ambient space is denoted by \mathbf{g} and the induced metric on the body in its reference configuration is denoted by $\mathbf{G} = \mathbf{g}|_{\mathcal{B}}$. A *deformation* is a map from \mathcal{B} to the ambient space, i.e., $\varphi : \mathcal{B} \rightarrow \mathcal{C} \subset \mathcal{S}$, where $\mathcal{C} = \varphi(\mathcal{B})$ is the current configuration. The tangent map of φ is the so-called *deformation gradient* $\mathbf{F} = T\varphi$ (a metric-independent map), which at each material point $X \in \mathcal{B}$ is a linear map $\mathbf{F}(X) : T_X\mathcal{B} \rightarrow T_{\varphi(X)}\mathcal{C}$. With respect to the coordinate charts $\{X^A\}$ and $\{x^a\}$ for \mathcal{B} and \mathcal{C} , respectively, the deformation gradient has components $F^a_A = \frac{\partial \varphi^a}{\partial X^A}$. The transpose of deformation gradient \mathbf{F}^T has components $(F^T)^A_a = g_{ab} F^b_B G^{AB}$. The *right Cauchy-Green strain tensor* is defined as $\mathbf{C} = \mathbf{F}^T \mathbf{F}$ with components $C^A_B = (F^T)^A_a F^a_B$. Thus, $C_{AB} = (g_{ab} \circ \varphi) F^a_A F^b_B$, which means that the right Cauchy-Green strain is the pull-back of the spatial metric to the reference configuration, i.e., $\mathbf{C}^b = \varphi^* \mathbf{g}$, where \flat is the flat operator induced by the metric \mathbf{G} (which lowers indices). The left Cauchy-Green strain is defined as $\mathbf{B}^\sharp = \varphi_*(\mathbf{g}^\sharp)$, which has components $B^{AB} = F^{-A}_a F^{-B}_b g^{ab}$, where F^{-A}_a are components of \mathbf{F}^{-1} . The spatial analogue of \mathbf{C}^b is defined as $\mathbf{c}^b = \varphi_* \mathbf{G}$, which has components $c_{ab} = F^{-A}_a F^{-B}_b G^{AB}$. Similarly, the spatial analogue of \mathbf{B}^\sharp is $\mathbf{b}^\sharp = \varphi_*(\mathbf{G}^\sharp)$, which has components $b^{ab} = F^a_A F^b_B G^{AB}$. Recall that $\mathbf{b} = \mathbf{c}^{-1}$. The two tensors \mathbf{C} and \mathbf{b} share the same principal invariants I_1 , I_2 , and I_3 , which are defined as [Ogden, 1984, Marsden and Hughes, 1994] $I_1 = \text{tr } \mathbf{b} = b^{ab} g_{ab}$, $I_2 = \frac{1}{2} (I_1^2 - \text{tr } \mathbf{b}^2) = \frac{1}{2} (I_1^2 - b^{ab} b^{cd} g_{ac} g_{bd})$, and $I_3 = \det \mathbf{b}$.

Implicit elasticity is defined by elastic materials that have the following implicit constitutive equation [Morgan, 1966, Rajagopal, 2003, 2007]¹

$$\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}. \quad (2.2)$$

Cauchy elastic and Green elastic (hyperelastic) solids are special cases of this class of elastic materials. Here, we confine ourselves to isotropic solids. In this case, the implicit constitutive equation can be rewritten as

¹In his study of *controllable states of stress*, Carroll [1973] considered a special subset of (2.2) constitutive equations in the form

$$\mathbf{b}^\sharp = \xi_0 \mathbf{g}^\sharp + \xi_1 \boldsymbol{\sigma} + \xi_2 \boldsymbol{\sigma}^2, \quad (2.1)$$

that he called an *invertible stress relation*. The response functions ξ_0 , ξ_1 , and ξ_2 are functions of the principal invariants of the Cauchy stress. According to Carroll [1973], a state of stress is controllable if it satisfies the equilibrium equations in the absence of body forces, and its corresponding strain is compatible for any response functions ξ_0 , ξ_1 , and ξ_2 . He proved that for isotropic elastic solids with constitutive equations of the form (2.1) controllable states of stress are homogeneous. He, however, noted that for a given elastic material with the constitutive equation (2.1), not all states of homogeneous stress are constitutively admissible.

[Rivlin and Ericksen, 1955, Morgan, 1966]²

$$\begin{aligned} & \beta_0 \mathbf{g}^\# + \beta_1 \boldsymbol{\sigma} + \beta_2 \boldsymbol{\sigma}^2 + \beta_3 \mathbf{b}^\# + \beta_4 \mathbf{c}^\# + \beta_5 (\boldsymbol{\sigma} \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}) + \beta_6 (\boldsymbol{\sigma}^2 \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}^2) \\ & + \beta_7 (\boldsymbol{\sigma} \mathbf{c}^\# + \mathbf{c}^\# \boldsymbol{\sigma}) + \beta_8 (\boldsymbol{\sigma}^2 \mathbf{c}^\# + \mathbf{c}^\# \boldsymbol{\sigma}^2) = \mathbf{0}, \end{aligned} \quad (2.4)$$

where β_i , $i = 0, \dots, 8$ are functions of the following ten invariants [Rivlin and Ericksen, 1955, Morgan, 1966, Rajagopal, 2003, 2007]:

$$\text{tr } \boldsymbol{\sigma}, \text{tr } \boldsymbol{\sigma}^2, \text{tr } \boldsymbol{\sigma}^3, \text{tr } \mathbf{b}^\#, \text{tr } \mathbf{b}^{2\#}, \text{tr } \mathbf{b}^{3\#}, \text{tr } (\boldsymbol{\sigma} \mathbf{b}^\#), \text{tr } (\boldsymbol{\sigma} \mathbf{b}^{2\#}), \text{tr } (\boldsymbol{\sigma}^2 \mathbf{b}^\#), \text{tr } (\boldsymbol{\sigma}^2 \mathbf{b}^{2\#}). \quad (2.5)$$

Equivalently, we can use the following invariants³

$$\begin{aligned} I_1 &= \text{tr } \boldsymbol{\sigma}, \quad I_2 = \text{tr } \boldsymbol{\sigma}^2, \quad I_3 = \text{tr } \boldsymbol{\sigma}^3, \quad I_4 = \text{tr } \mathbf{b}^\#, \quad I_5 = \frac{1}{2} [I_4^2 - \text{tr } \mathbf{b}^{2\#}], \quad I_6 = \det \mathbf{b}^\#, \\ I_7 &= \text{tr } (\boldsymbol{\sigma} \mathbf{b}^\#), \quad I_8 = \text{tr } (\boldsymbol{\sigma} \mathbf{b}^{2\#}), \quad I_9 = \text{tr } (\boldsymbol{\sigma}^2 \mathbf{b}^\#), \quad I_{10} = \text{tr } (\boldsymbol{\sigma}^2 \mathbf{b}^{2\#}). \end{aligned} \quad (2.6)$$

Hence, the implicit constitutive equations are rewritten as⁴

$$\begin{aligned} \mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) &= \alpha_0 \mathbf{g}^\# + \alpha_1 \boldsymbol{\sigma} + \alpha_2 \boldsymbol{\sigma}^2 + \alpha_3 \mathbf{b}^\# + \alpha_4 \mathbf{c}^\# + \alpha_5 (\boldsymbol{\sigma} \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}) + \alpha_6 (\boldsymbol{\sigma}^2 \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}^2) \\ &+ \alpha_7 (\boldsymbol{\sigma} \mathbf{c}^\# + \mathbf{c}^\# \boldsymbol{\sigma}) + \alpha_8 (\boldsymbol{\sigma}^2 \mathbf{c}^\# + \mathbf{c}^\# \boldsymbol{\sigma}^2) = \mathbf{0}, \end{aligned} \quad (2.7)$$

where $\alpha_i = \alpha_i(I_1, \dots, I_{10})$, $i = 0, \dots, 8$. Note that with respect to a coordinate chart $\{x^a\}$, (2.7) is written as $(a, b = 1, 2, 3)$

$$\begin{aligned} & \alpha_0 g^{ab} + \alpha_1 \sigma^{ab} + \alpha_2 \sigma^{2ab} + \alpha_3 b^{ab} + \alpha_4 c^{ab} + \alpha_5 (\sigma^{am} b_m^b + b_m^a \sigma^{mb}) + \alpha_6 (\sigma^{2am} b_m^b + b_m^a \sigma^{2mb}) \\ & + \alpha_7 (\sigma^{am} c_m^b + c_m^a \sigma^{mb}) + \alpha_8 (\sigma^{2am} c_m^b + c_m^a \sigma^{2mb}) = 0, \end{aligned} \quad (2.8)$$

where $\sigma^{2ab} = \sigma^{am} g_{mn} \sigma^{nb}$.

Definition 2.1. The *implicit-elasticity stress-strain space* \mathbb{S} is a twelve-dimensional submanifold of \mathbb{R}^{12} defined as

$$\mathbb{S} = \{s \in \mathbb{R}^{12} : s = (b^{ab}, \sigma^{cd}), a \leq b = 1, 2, 3, c \leq d = 1, 2, 3, \mathbf{b} \text{ positive-definite and compatible}\}. \quad (2.9)$$

The *implicit-elasticity stress-strain manifold* \mathfrak{S} of an implicit-elastic material is the six-dimensional submanifold of \mathbb{S} defined by the six equations $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$.

Definition 2.2. In implicit elasticity, a homogeneous deformation (that has a constant \mathbf{b}) is *constitutively admissible* if $(\mathbf{b}, \boldsymbol{\sigma}) \in \mathfrak{S}$, i.e., if there exists $\boldsymbol{\sigma}$ such that $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$.

Example 2.3. In (2.7) suppose $\alpha_5 = \alpha_6 = \alpha_7 = \alpha_8 = 0$, i.e.,

$$\alpha_2 \boldsymbol{\sigma}^2 + \alpha_1 \boldsymbol{\sigma} = -\alpha_0 \mathbf{g}^\# - \alpha_3 \mathbf{b}^\# - \alpha_4 \mathbf{c}^\#. \quad (2.10)$$

We denote the eigenvalues of $\boldsymbol{\sigma}$ and $\mathbf{b}^\#$ by σ_i and λ_i^2 , $i = 1, 2, 3$, respectively. For this class of materials, $\boldsymbol{\sigma}$ and $\mathbf{b}^\#$ have the same eigenvectors and can be diagonalized simultaneously.⁵ Therefore, with respect to the eigenbasis one has

$$\alpha_2 \sigma_i^2 + \alpha_1 \sigma_i + (\alpha_0 + \alpha_3 \lambda_i^2 + \alpha_4 \lambda_i^{-2}) = 0, \quad i = 1, 2, 3. \quad (2.11)$$

²In [Rivlin and Ericksen, 1955, Morgan, 1966], this is written as

$$\begin{aligned} & \beta_0 \mathbf{g}^\# + \beta_1 \boldsymbol{\sigma} + \beta_2 \boldsymbol{\sigma}^2 + \beta_3 \mathbf{b}^\# + \beta_4 \mathbf{b}^{2\#} + \beta_5 (\boldsymbol{\sigma} \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}) + \beta_6 (\boldsymbol{\sigma}^2 \mathbf{b}^\# + \mathbf{b}^\# \boldsymbol{\sigma}^2) \\ & + \beta_7 (\boldsymbol{\sigma} \mathbf{b}^{2\#} + \mathbf{b}^{2\#} \boldsymbol{\sigma}) + \beta_8 (\boldsymbol{\sigma}^2 \mathbf{b}^{2\#} + \mathbf{b}^{2\#} \boldsymbol{\sigma}^2) = \mathbf{0}. \end{aligned} \quad (2.3)$$

However, from the Cayley-Hamilton theorem we know that $\mathbf{b}^{2\#}$ is functionally dependent on $\mathbf{b}^\#$ and $\mathbf{b}^{-\#} = \mathbf{c}^\#$.

³Clearly, one can use the pair I_4 and I_5 instead of I_4 and $\text{tr } \mathbf{b}^{2\#}$. Using the Cayley-Hamilton theorem, one can show that $\text{tr } \mathbf{b}^{3\#} = I_4^3 - 3I_4 I_5 + 3I_6$ [Spencer, 1971].

⁴Assuming that the initial configuration is stress free, $\alpha_0 + \alpha_3 + \alpha_4 = 0$ when $(I_1, \dots, I_{10}) = (0, 0, 0, 3, 3, 1, 0, 0, 0, 0)$.

⁵It should be emphasized that for the more general class of elastic solids with constitutive equations (2.7), $\boldsymbol{\sigma}$ and $\mathbf{b}^\#$ do not have the same eigenvectors, in general.

These quadratic equations (in σ_i) have real solutions if and only if $\alpha_1^2 - 4\alpha_2(\alpha_0 + \alpha_3\lambda_i^2 + \alpha_4\lambda_i^{-2}) \geq 0$, $i = 1, 2, 3$. Thus, \mathbf{b}^\sharp is constitutively inadmissible if for at least one value of i one has

$$\alpha_1^2 - 4\alpha_2(\alpha_0 + \alpha_3\lambda_i^2 + \alpha_4\lambda_i^{-2}) < 0. \quad (2.12)$$

We observe that not every homogeneous deformation is constitutively admissible.

Example 2.4. Consider the particular case of (2.7) for which $\alpha_1 = \alpha_2 = \alpha_6 = \alpha_7 = \alpha_8 = 0$ and $\alpha_5 \neq 0$, i.e.,

$$\boldsymbol{\sigma}\mathbf{b}^\sharp + \mathbf{b}^\sharp\boldsymbol{\sigma} = \eta_0\mathbf{g}^\sharp + \eta_1\mathbf{b}^\sharp + \eta_2\mathbf{c}^\sharp. \quad (2.13)$$

For a homogeneous deformation and constant η_0, η_1, η_2 (these are constant when stress is homogeneous as well), (2.13) is a Sylvester equation [Sylvester, 1884, Bhatia and Rosenthal, 1997, Gantmakher, 2000]. There is a unique solution for stress if and only if \mathbf{b}^\sharp and $-\mathbf{b}^\sharp$ do not share any eigenvalues. This is the case because all the eigenvalues of \mathbf{b}^\sharp are positive and consequently those of $-\mathbf{b}^\sharp$ are all negative. Therefore, any homogeneous \mathbf{b}^\sharp has a unique corresponding homogeneous stress.

3 Universal Deformations in Compressible Isotropic Implicit Elasticity

Universal deformations for Cauchy-elastic solids. Before considering the general case of implicit elasticity, we start by characterizing the universal deformations of isotropic Cauchy-elastic solids. This class of materials is defined by an explicit constitutive equations of the form:

$$\boldsymbol{\sigma} = \beta_0\mathbf{g}^\sharp + \beta_1\mathbf{b}^\sharp + \beta_2\mathbf{c}^\sharp, \quad (3.1)$$

where $\beta_i = \beta_i(I_1, I_2, I_3)$, $i = 1, 2, 3$. Equilibrium equations in the absence of body forces are written as (note that metric is covariantly constant, and hence, $\text{div } \mathbf{g}^\sharp = \mathbf{0}$)

$$\mathbf{0} = \text{div } \boldsymbol{\sigma} = \text{div } [\beta_0\mathbf{g}^\sharp + \beta_1\mathbf{b}^\sharp + \beta_2\mathbf{c}^\sharp] = \beta_1 \text{div } \mathbf{b}^\sharp + \beta_2 \text{div } \mathbf{c}^\sharp + \nabla\beta_0 + \mathbf{b}^\sharp \cdot \nabla\beta_1 + \mathbf{c}^\sharp \cdot \nabla\beta_2. \quad (3.2)$$

Then, it is noted that

$$\nabla\beta_i = \sum_{j=1}^3 \beta_{ij} \nabla I_j, \quad \beta_{ij} = \frac{\partial \beta_i}{\partial I_j}, \quad i = 1, 2, 3. \quad (3.3)$$

Thus, we have

$$\beta_1 \text{div } \mathbf{b}^\sharp + \beta_2 \text{div } \mathbf{c}^\sharp + \sum_{j=1}^3 \beta_{0j} \nabla I_j + \sum_{j=1}^3 \beta_{1j} \mathbf{b}^\sharp \cdot \nabla I_j + \sum_{j=1}^3 \beta_{2j} \mathbf{c}^\sharp \cdot \nabla I_j = \mathbf{0}. \quad (3.4)$$

This identity must hold for arbitrary response functions β_0 , β_1 , and β_2 . Knowing that any derivative of a response function is functionally independent from the response functions and all the other derivatives, one concludes that in (3.4) the coefficient of each response function and its partial derivatives must be identically zero. This immediately implies that [Yavari, 2024]

$$\text{div } \mathbf{b}^\sharp = \text{div } \mathbf{c}^\sharp = \mathbf{0}, \quad \text{and} \quad \text{grad } I_j = \mathbf{0}, \quad i = 1, 2, 3. \quad (3.5)$$

These constraints are identical to the universality constraints of isotropic hyperelasticity and imply that the universal deformations are homogeneous [Ericksen, 1955]. It should be emphasized that universal deformations are independent of the response functions β_0 , β_1 , and β_2 . However, for a homogeneous body with a given triplet of response functions $(\beta_0, \beta_1, \beta_2)$, stress explicitly depends on the response functions, but is nevertheless uniform.

Equilibrium stress with constant principal invariants is homogeneous. To extend this analysis to the more general case of isotropic implicit elasticity given by constitutive equations (2.7) we will need the following general result.

Lemma 3.1. *If $\boldsymbol{\sigma}$ is an equilibrium stress, i.e. $\operatorname{div} \boldsymbol{\sigma} = \mathbf{0}$, with constant principal invariants, then it is homogeneous.*

Proof. The Cauchy stress can be written in the following spectral representation [Ogden, 1984]:

$$\boldsymbol{\sigma} = \sigma_1 \mathbf{a} \otimes \mathbf{a} + \sigma_2 \mathbf{b} \otimes \mathbf{b} + \sigma_3 \mathbf{c} \otimes \mathbf{c}, \quad (3.6)$$

where $\sigma_1, \sigma_2, \sigma_3$ and $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are the principal stresses and their corresponding principal directions. Note that $\mathbf{a} \otimes \mathbf{a} + \mathbf{b} \otimes \mathbf{b} + \mathbf{c} \otimes \mathbf{c} = \mathbf{g}^\sharp$. Thus, the Cauchy stress can be written

$$\boldsymbol{\sigma} = (\sigma_1 - \sigma_3) \mathbf{a} \otimes \mathbf{a} + (\sigma_2 - \sigma_3) \mathbf{b} \otimes \mathbf{b} + \sigma_3 \mathbf{g}^\sharp. \quad (3.7)$$

If it has constant invariants then it has constant eigenvalues. Moreover, being an equilibrium stress, it is divergence-free. Therefore, there are three possibilities for the eigenvalues: (i) $\sigma_1 = \sigma_2 = \sigma_3$, (ii) $\sigma_2 = \sigma_3 \neq \sigma_1$, and (iii) the three eigenvalues are distinct.

Case (i): When the three eigenvalues of $\boldsymbol{\sigma}$ are equal, we have $\boldsymbol{\sigma} = \sigma_1 \mathbf{g}^\sharp$, which is a covariantly constant tensor, i.e., a homogeneous tensor in the Euclidean space.

Case (ii): when $\sigma_2 = \sigma_3 \neq \sigma_1$, the stress has the following representation

$$\boldsymbol{\sigma} = (\sigma_1 - \sigma_3) \mathbf{a} \otimes \mathbf{a} + \sigma_3 \mathbf{g}^\sharp. \quad (3.8)$$

Thus, $\operatorname{div} \boldsymbol{\sigma} = (\sigma_1 - \sigma_3) \operatorname{div}(\mathbf{a} \otimes \mathbf{a}) = \mathbf{0}$, and hence $\operatorname{div}(\mathbf{a} \otimes \mathbf{a}) = \mathbf{0}$. Note that

$$\operatorname{div}(\mathbf{a} \otimes \mathbf{a}) = \nabla_{\mathbf{a}} \mathbf{a} + (\operatorname{div} \mathbf{a}) \mathbf{a}, \quad \text{or in components} \quad (\mathbf{a}^m \mathbf{a}^n)_{|n} = \mathbf{a}^m_{|n} \mathbf{a}^n + \mathbf{a}^n_{|n} \mathbf{a}^m. \quad (3.9)$$

Therefore, equilibrium dictates that

$$\nabla_{\mathbf{a}} \mathbf{a} + (\operatorname{div} \mathbf{a}) \mathbf{a} = \mathbf{0}, \quad \text{or in components} \quad \mathbf{a}^m_{|n} \mathbf{a}^n + \mathbf{a}^n_{|n} \mathbf{a}^m = 0. \quad (3.10)$$

Knowing that $\mathbf{a}^m \mathbf{a}_m = 1$, one has $0 = (\mathbf{a}^m \mathbf{a}_m)_{|b} = \mathbf{a}^m_{|b} \mathbf{a}_m + \mathbf{a}^m \mathbf{a}_{m|b} = \mathbf{a}^m_{|b} \mathbf{a}_m + \mathbf{a}_m \mathbf{a}^m_{|b} = 2\mathbf{a}^m_{|b} \mathbf{a}_m$. Thus, for any unit vector \mathbf{a}

$$\mathbf{a}_m \mathbf{a}^m_{|b} = 0. \quad (3.11)$$

Taking the dot product of the vector equation (3.10) with \mathbf{a} and using (3.11), one obtains

$$0 = \mathbf{a}_m \mathbf{a}^m_{|n} \mathbf{a}^n + \mathbf{a}^n_{|n} \mathbf{a}^m \mathbf{a}_m = \mathbf{a}^n_{|n}. \quad (3.12)$$

Thus, we have

$$\operatorname{div} \mathbf{a} = \mathbf{0}. \quad (3.13)$$

Using (3.13) in (3.10), one concludes that

$$\nabla_{\mathbf{a}} \mathbf{a} = \mathbf{0}. \quad (3.14)$$

This is the geodesic equation that tells us that the integral curves of \mathbf{a} are geodesics. But we know that in the Euclidean space geodesics are straight lines. This in turn implies that the stress is homogeneous.

Case (iii): The three eigenvalues are distinct. In this case, the stress has the following representation (3.7) and we have

$$\operatorname{div} \boldsymbol{\sigma} = (\sigma_1 - \sigma_3) \nabla_{\mathbf{a}} \mathbf{a} + (\sigma_2 - \sigma_3) \nabla_{\mathbf{b}} \mathbf{b} = \mathbf{0}. \quad (3.15)$$

In components

$$(\sigma_1 - \sigma_3) \mathbf{a}^m_{|n} \mathbf{a}^n + (\sigma_2 - \sigma_3) \mathbf{b}^m_{|n} \mathbf{b}^n = 0. \quad (3.16)$$

Hence, by taking the dot product of both sides of the above identity with \mathbf{a} , one obtains

$$0 = (\sigma_1 - \sigma_3) \mathbf{a}_m \mathbf{a}^m_{|n} \mathbf{a}^n + (\sigma_2 - \sigma_3) \mathbf{a}_m \mathbf{b}^m_{|n} \mathbf{b}^n = (\sigma_2 - \sigma_3) \mathbf{a}_m \mathbf{b}^m_{|n} \mathbf{b}^n = 0, \quad (3.17)$$

and hence

$$\mathbf{a}_m \mathbf{b}^m|_n \mathbf{b}^n = 0 \quad \text{or} \quad \mathbf{a} \cdot \nabla_{\mathbf{b}} \mathbf{b} = 0. \quad (3.18)$$

Similarly

$$\mathbf{b} \cdot \nabla_{\mathbf{a}} \mathbf{a} = 0. \quad (3.19)$$

In summary, $\mathbf{a} \cdot \nabla_{\mathbf{a}} \mathbf{a} = \mathbf{b} \cdot \nabla_{\mathbf{a}} \mathbf{a} = \mathbf{a} \cdot \nabla_{\mathbf{b}} \mathbf{b} = \mathbf{b} \cdot \nabla_{\mathbf{b}} \mathbf{b} = 0$. Therefore

$$\nabla_{\mathbf{a}} \mathbf{a} = k_1 \mathbf{c}, \quad \nabla_{\mathbf{b}} \mathbf{b} = k_2 \mathbf{c}. \quad (3.20)$$

Instead of the spectral representation (3.7) one can equivalently write

$$\boldsymbol{\sigma} = (\sigma_1 - \sigma_2) \mathbf{a} \otimes \mathbf{a} + (\sigma_3 - \sigma_2) \mathbf{c} \otimes \mathbf{c} + \sigma_2 \mathbf{g}^\sharp. \quad (3.21)$$

Using this representation and following a similar argument that led to the identity (3.19), one can show that $\mathbf{c} \cdot \nabla_{\mathbf{a}} \mathbf{a} = 0$, and hence $k_1 = 0$, i.e., $\nabla_{\mathbf{a}} \mathbf{a} = \mathbf{0}$. One can also use the following spectral representation

$$\boldsymbol{\sigma} = (\sigma_2 - \sigma_1) \mathbf{b} \otimes \mathbf{b} + (\sigma_3 - \sigma_1) \mathbf{c} \otimes \mathbf{c} + \sigma_1 \mathbf{g}^\sharp. \quad (3.22)$$

Using this representation and following a similar argument that led to the identity (3.18), one can show that $\mathbf{c} \cdot \nabla_{\mathbf{b}} \mathbf{b} = 0$, and hence $k_2 = 0$, i.e., $\nabla_{\mathbf{b}} \mathbf{b} = \mathbf{0}$. From $\nabla_{\mathbf{a}} \mathbf{a} = \nabla_{\mathbf{b}} \mathbf{b} = \mathbf{0}$ we conclude that the integral curves of \mathbf{a} and \mathbf{b} are orthogonal straight lines. This implies that the integral curves of \mathbf{c} are straight lines as well and orthogonal to those of \mathbf{a} and \mathbf{b} . This again implies that the Cauchy stress is homogeneous. \square

Universal deformations for a special class of implicit-elastic solids. Next, we consider the special subclass of materials with constitutive equations (2.1), i.e.,

$$\mathbf{b}^\sharp = \xi_0 \mathbf{g}^\sharp + \xi_1 \boldsymbol{\sigma} + \xi_2 \boldsymbol{\sigma}^2. \quad (3.23)$$

We take the divergence of both sides and use the fact that $\operatorname{div} \boldsymbol{\sigma} = \mathbf{0}$ (equilibrium equations in the absence of body forces) and $\operatorname{div} \mathbf{g}^\sharp = \mathbf{0}$ (metric is covariantly constant):

$$\nabla \xi_0 + \boldsymbol{\sigma} \nabla \xi_1 + \boldsymbol{\sigma}^2 \nabla \xi_2 + \xi_2 \operatorname{div} \boldsymbol{\sigma}^2 - \operatorname{div} \mathbf{b}^\sharp = \mathbf{0}. \quad (3.24)$$

The above identity holds for arbitrary response functions ξ_0, ξ_1, ξ_2 . Let us replace ξ_0 by $\zeta \xi_0$:⁶ $\zeta \nabla \xi_0 + \boldsymbol{\sigma} \nabla \xi_1 + \boldsymbol{\sigma}^2 \nabla \xi_2 + \xi_2 \operatorname{div} \boldsymbol{\sigma}^2 - \operatorname{div} \mathbf{b}^\sharp = \mathbf{0}$. Differentiating both sides of the above identity with respect to ζ one obtains $\nabla \xi_0 = \mathbf{0}$, and hence the principal invariants of Cauchy stress are constant. Thus, Cauchy stress is homogeneous and ξ_0, ξ_1, ξ_2 are constant. Hence, \mathbf{b}^\sharp is homogeneous. Therefore, universal deformations are homogeneous. In Example 2.3 we saw that not every homogeneous deformation is constitutively admissible for this class of elastic materials. In other words, unlike Cauchy (and Green) elasticity, the set of universal deformations depends on the response functions. All we know is that for any material within this class the set of universal deformations is a subset of homogeneous deformations.

Remark 3.2. To prove that universal deformations in homogeneous compressible isotropic hyperelastic solids are homogeneous, Ericksen [1955] used the equilibrium equations in the absence of body forces, the necessity for these equations to hold for any energy function, and the compatibility equations of the left Cauchy-Green strain. A simpler proof was presented in [Saccomandi, 2001b] by employing the following two specific energy functions: $W(I_1, I_2, I_3) = \mu_1 I_1 + \mu_2 I_2 + \mu_3 I_3$ and $\hat{W}(I_1, I_2, I_3) = \hat{\mu}_1 I_1^2 + \hat{\mu}_2 I_2^2 + \hat{\mu}_3 I_3^2$, where μ_i and $\hat{\mu}_i$, $i = 1, 2, 3$, are arbitrary constants. Casey [2004] demonstrated that Ericksen's result can be proved using the simpler energy functions $W = \mu_1 I_1$ and $\hat{W} = \hat{\mu}_1 I_1^2$. Furthermore, he concluded that universal deformations for “all homogeneous elastic materials” must be homogeneous. While this is true, it is important to note that, in general, not every homogeneous deformation is constitutively admissible for a given elastic material, as demonstrated in Example 2.3.

⁶ ζ is an arbitrary scaling factor, i.e., $\zeta \in \mathbb{R}$

Universal deformations in implicit-elastic solids. Universal deformations correspond to those deformations that satisfy the equilibrium equations in the absence of body forces for arbitrary response functions α_i , $i = 0, \dots, 8$. We pick an arbitrary deformation, which has a left Cauchy-Green strain \mathbf{b}^\sharp (and its corresponding \mathbf{c}^\sharp). The implicit constitutive equation (2.7) determines the corresponding stress $\boldsymbol{\sigma}$ (that may not be unique) if \mathbf{b}^\sharp is constitutively admissible. Note that for this fixed deformation, $\boldsymbol{\sigma} = \boldsymbol{\sigma}(\alpha_0, \dots, \alpha_8)$. This means that when we choose different materials within the class of materials (2.7), $\boldsymbol{\sigma}$ changes. In other words, a given deformation has, in general, infinitely many corresponding stresses within the material class. Any stress $\boldsymbol{\sigma}$ corresponding to the fixed \mathbf{b}^\sharp and \mathbf{c}^\sharp must satisfy (2.7) everywhere in the body. We take the (spatial) divergence of both sides of (2.7) and note that $\text{div } \boldsymbol{\sigma} = \mathbf{0}$ (equilibrium equations in the absence of body forces) and $\text{div } \mathbf{g}^\sharp = \mathbf{0}$ (metric is covariantly constant):

$$\begin{aligned} & \nabla \alpha_0 + \boldsymbol{\sigma} \cdot \nabla \alpha_1 + \boldsymbol{\sigma}^2 \cdot \nabla \alpha_2 + \alpha_2 \text{div } \boldsymbol{\sigma}^2 + \alpha_3 \text{div } \mathbf{b}^\sharp + \mathbf{b}^\sharp \cdot \nabla \alpha_3 + \alpha_4 \text{div } \mathbf{c}^\sharp + \mathbf{c}^\sharp \cdot \nabla \alpha_4 \\ & + \alpha_5 [\nabla \boldsymbol{\sigma} \cdot \mathbf{b}^\sharp + \nabla \mathbf{b}^\sharp \cdot \boldsymbol{\sigma} + \boldsymbol{\sigma} \text{div } \mathbf{b}^\sharp] + (\boldsymbol{\sigma} \mathbf{b}^\sharp + \mathbf{b}^\sharp \boldsymbol{\sigma}) \cdot \nabla \alpha_5 \\ & + \alpha_6 [\nabla \boldsymbol{\sigma}^2 \cdot \mathbf{b}^\sharp + \nabla \mathbf{b}^\sharp \cdot \boldsymbol{\sigma}^2 + \boldsymbol{\sigma}^2 \text{div } \mathbf{b}^\sharp + \mathbf{b}^\sharp \text{div } \boldsymbol{\sigma}^2] + (\boldsymbol{\sigma}^2 \mathbf{b}^\sharp + \mathbf{b}^\sharp \boldsymbol{\sigma}^2) \cdot \nabla \alpha_6 \\ & + \alpha_7 [\nabla \boldsymbol{\sigma} \cdot \mathbf{c}^\sharp + \nabla \mathbf{c}^\sharp \cdot \boldsymbol{\sigma} + \boldsymbol{\sigma} \text{div } \mathbf{c}^\sharp] + (\boldsymbol{\sigma} \mathbf{c}^\sharp + \mathbf{c}^\sharp \boldsymbol{\sigma}) \cdot \nabla \alpha_7 \\ & + \alpha_8 [\nabla \boldsymbol{\sigma}^2 \cdot \mathbf{c}^\sharp + \nabla \mathbf{c}^\sharp \cdot \boldsymbol{\sigma}^2 + \boldsymbol{\sigma}^2 \text{div } \mathbf{c}^\sharp + \mathbf{c}^\sharp \text{div } \boldsymbol{\sigma}^2] + (\boldsymbol{\sigma}^2 \mathbf{c}^\sharp + \mathbf{c}^\sharp \boldsymbol{\sigma}^2) \cdot \nabla \alpha_8 = \mathbf{0}. \end{aligned} \quad (3.25)$$

For fixed \mathbf{b}^\sharp and \mathbf{c}^\sharp , $\boldsymbol{\sigma} = \boldsymbol{\sigma}(\alpha_0, \dots, \alpha_8)$, and the above identity must hold everywhere in the deformed body for arbitrary response functions $\alpha_i = \alpha_i(I_1, \dots, I_{10})$, $i = 0, \dots, 8$. The response functions and their derivatives can vary independently of each other. Specifically, $\nabla \alpha_0$ can vary independently from α_0 , α_j ($j = 1, \dots, 8$), and $\nabla \alpha_j$ ($j = 1, \dots, 8$). Note that $\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}^2$ do not depend on the derivative of α_0 with respect to any of the ten invariants. Therefore, one concludes that

$$\nabla \alpha_0 = \sum_{j=1}^{10} \frac{\partial \alpha_0}{\partial I_j} \nabla I_j = \mathbf{0}. \quad (3.26)$$

Therefore, the arbitrariness of the derivatives of α_0 with respect to the ten invariants implies that

$$I_1, \dots, I_{10} \text{ are constant}. \quad (3.27)$$

I_1, I_2, I_3 being constant tells us that the Cauchy stress is homogeneous. Thus, (3.25) is simplified to read

$$\begin{aligned} & \alpha_3 \text{div } \mathbf{b}^\sharp + \alpha_4 \text{div } \mathbf{c}^\sharp + \alpha_5 \boldsymbol{\sigma} \text{div } \mathbf{b}^\sharp + \alpha_6 [\nabla \mathbf{b}^\sharp \cdot \boldsymbol{\sigma}^2 + \boldsymbol{\sigma}^2 \text{div } \mathbf{b}^\sharp] \\ & + \alpha_7 [\nabla \mathbf{c}^\sharp \cdot \boldsymbol{\sigma} + \boldsymbol{\sigma} \text{div } \mathbf{c}^\sharp] + \alpha_8 [\nabla \mathbf{c}^\sharp \cdot \boldsymbol{\sigma}^2 + \boldsymbol{\sigma}^2 \text{div } \mathbf{c}^\sharp] = \mathbf{0}. \end{aligned} \quad (3.28)$$

In the constitutive equation (2.7) let us assume that α_0 , α_1 , α_3 , and α_4 are the only nonzero response functions (this corresponds to Cauchy elasticity). Thus, in the above identity the response function α_3 and α_4 can be chosen arbitrarily, and hence $\text{div } \mathbf{b}^\sharp = \text{div } \mathbf{c}^\sharp = \mathbf{0}$. We observe that the universality constraints of implicit elasticity include those of Cauchy elasticity, i.e., (3.5). Therefore, universal deformations are homogeneous, and (3.28) is now trivially satisfied. Therefore we have proved the following result.

Proposition 3.3. *In compressible isotropic implicit elasticity, all universal deformations are homogeneous.*

Remark 3.4. It should be emphasized that for a given implicit-elastic material not every homogeneous deformation is constitutively admissible. In other words, the set of universal deformations is material dependent, in general. All one knows is that for a given material within the class of implicit-elastic materials, the set of universal deformations is a subset of homogeneous deformations. This suggests that homogeneous deformations remain useful for characterizing material properties of implicit-elastic materials.

4 Conclusions

Universal deformations have been extensively studied for compressible and incompressible hyperelastic solids. In recent years, the study of universal deformations has been expanded to encompass anelasticity, as well

as anisotropic and inhomogeneous hyperelastic bodies. More recently, universal deformations have been studied for compressible and incompressible Cauchy elasticity, which includes hyperelasticity as a subset. It is known that there are larger classes of elastic solids that include, for instance, Cauchy elasticity. One such class of solids was introduced by Morgan [1966] characterized by implicit constitutive equations of the form $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$. The study of this class of elastic solids has been revived in the past twenty years by Rajagopal and his collaborators [Rajagopal, 2003, 2007, Bustamante, 2009, Bustamante and Rajagopal, 2011]. In this paper we investigated the problem of finding the universal deformations of compressible isotropic solids with the implicit constitutive equations $\mathbf{f}(\boldsymbol{\sigma}, \mathbf{b}) = \mathbf{0}$. We showed that the universal deformations for these materials are homogeneous. However, we observed that, in general, not every homogeneous deformation is constitutively admissible for a given implicit-elastic material. This implies that in implicit elasticity the set of universal deformations is material dependent. However, for any given implicit-elastic material, the set of universal deformations is a subset of the set of homogeneous deformations. This suggests that homogenous deformations can still be used to characterize the material properties of such solids.

Acknowledgments

This work was supported by NSF – Grant No. CMMI 1939901.

References

- R. Bhatia and P. Rosenthal. How and why to solve the operator equation $ax - xb = y$. *Bulletin of the London Mathematical Society*, 29(1):1–21, 1997.
- R. Bustamante. Some topics on a new class of elastic bodies. *Proceedings of the Royal Society A*, 465(2105):1377–1392, 2009.
- R. Bustamante. Some universal solutions for a class of incompressible elastic body that is not green elastic: The case of large elastic deformations. *The Quarterly Journal of Mechanics and Applied Mathematics*, 73(2):177–199, 2020a.
- R. Bustamante. Some universal solutions for incompressible elastic bodies that are not Green elastic. *International Journal of Engineering Science*, 149:103223, 2020b.
- R. Bustamante and K. Rajagopal. Solutions of some simple boundary value problems within the context of a new class of elastic materials. *International Journal of Non-Linear Mechanics*, 46(2):376–386, 2011.
- M. M. Carroll. Controllable states of stress for compressible elastic solids. *Journal of Elasticity*, 3:57–61, 1973.
- J. Casey. On Ericksen’s theorem for unconstrained hyperelastic materials. *Journal of Elasticity*, 76:191–197, 2004.
- H. Chi, C. Talischi, O. Lopez-Pamies, and G. H. Paulino. Polygonal finite elements for finite elasticity. *International Journal for Numerical Methods in Engineering*, 101(4):305–328, 2015.
- R. De Pascalis, K. Rajagopal, and G. Saccomandi. Remarks on the use and misuse of the semi-inverse method in the nonlinear theory of elasticity. *Quarterly journal of Mechanics and Applied Mathematics*, 62(4):451–464, 2009.
- T. Doyle and J. L. Ericksen. Nonlinear elasticity. *Advances in Applied Mechanics*, 4:53–115, 1956.
- E. Dragoni. The radial compaction of a hyperelastic tube as a benchmark in compressible finite elasticity. *International Journal of Non-Linear Mechanics*, 31(4):483–493, 1996.
- J. L. Ericksen. Deformations possible in every isotropic, incompressible, perfectly elastic body. *Zeitschrift für Angewandte Mathematik und Physik*, 5(6):466–489, 1954.

- J. L. Ericksen. Deformations possible in every compressible, isotropic, perfectly elastic material. *Studies in Applied Mathematics*, 34(1-4):126–128, 1955.
- J. L. Ericksen and R. S. Rivlin. Large elastic deformations of homogeneous anisotropic materials. *Journal of Rational Mechanics and Analysis*, 3:281–301, 1954.
- R. L. Fosdick. Remarks on Compatibility. *Modern Developments in the Mechanics of Continua*, pages 109–127, 1966.
- B. Gairola. *Nonlinear elastic problems*. F.R.N. Nabarro (Ed.), Dislocations in Solids. North-Holland Publishing Co., Amsterdam, 1979.
- F. R. Gantmakher. *The Theory of Matrices*, volume 131. American Mathematical Soc., 2000.
- A. Golgoon and A. Yavari. Nonlinear elastic inclusions in anisotropic solids. *Journal of Elasticity*, 130(2): 239–269, 2018a.
- A. Golgoon and A. Yavari. Line and point defects in nonlinear anisotropic solids. *Zeitschrift für angewandte Mathematik und Physik*, 69:1–28, 2018b.
- A. Golgoon and A. Yavari. On Hashin’s hollow cylinder and sphere assemblages in anisotropic nonlinear elasticity. *Journal of Elasticity*, 146(1):65–82, 2021.
- C. Goodbrake, A. Yavari, and A. Goriely. The anelastic Ericksen problem: Universal deformations and universal eigenstrains in incompressible nonlinear anelasticity. *Journal of Elasticity*, 142(2):291–381, 2020.
- A. Goriely. *The Mathematics and Mechanics of Biological Growth*, volume 45. Springer, 2017.
- M. E. Gurtin. The linear theory of elasticity. In *Handbuch der Physik, Band VIa/2*. Springer-Verlag, Berlin, 1972.
- Z. Hashin. Large isotropic elastic deformation of composites and porous media. *International Journal of Solids and Structures*, 21(7):711–720, 1985.
- W. W. Klingbeil and R. T. Shield. On a class of solutions in plane finite elasticity. *Zeitschrift für angewandte Mathematik und Physik*, 17(4):489–511, 1966.
- J. K. Knowles. Universal states of finite anti-plane shear: Ericksen’s problem in miniature. *The American Mathematical Monthly*, 86(2):109–113, 1979.
- A. Kumar and A. Yavari. Nonlinear mechanics of remodeling. *Journal of the Mechanics and Physics of Solids*, 181:105449, 2023.
- V. Lee and K. Bhattacharya. Universal deformations of incompressible nonlinear elasticity as applied to ideal liquid crystal elastomers. *Journal of Elasticity*, pages 1–27, 2023.
- O. Lopez-Pamies, J. Moraleda, J. Segurado, and J. Llorca. On the extremal properties of Hashin’s hollow cylinder assemblage in nonlinear elasticity. *Journal of Elasticity*, 107:1–10, 2012.
- J. E. Marsden and T. J. R. Hughes. *Mathematical Foundations of Elasticity*. Dover Publications, New York, 1994.
- L. A. Mihai and A. Goriely. Controllable deformations of unconstrained ideal nematic elastomers. *Journal of Elasticity*, pages 1–12, 2023.
- A. J. A. Morgan. Some properties of media defined by constitutive equations in implicit form. *International Journal of Engineering Science*, 4(2):155–178, 1966.
- R. W. Ogden. *Non-Linear Elastic Deformations*. Dover, 1984.

- D. A. Polignone and C. O. Horgan. Pure torsion of compressible non-linearly elastic circular cylinders. *Quarterly of Applied Mathematics*, 49(3):591–607, 1991.
- S. P. Pradhan and A. Yavari. Accretion–ablation mechanics. *Philosophical Transactions of the Royal Society A*, 381(2263):20220373, 2023.
- K. R. Rajagopal. On implicit constitutive theories. *Applications of Mathematics*, 48:279–319, 2003.
- K. R. Rajagopal. The elasticity of elasticity. *Zeitschrift für angewandte Mathematik und Physik*, 58:309–317, 2007.
- R. S. Rivlin. Large elastic deformations of isotropic materials IV. Further developments of the general theory. *Philosophical Transactions of the Royal Society of London A*, 241(835):379–397, 1948.
- R. S. Rivlin. Large elastic deformations of isotropic materials. V. The problem of flexure. *Proceedings of the Royal Society of London A*, 195(1043):463–473, 1949a.
- R. S. Rivlin. A note on the torsion of an incompressible highly elastic cylinder. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 45, pages 485–487. Cambridge University Press, 1949b.
- R. S. Rivlin and J. L. Ericksen. Stress-deformation relations for isotropic materials. *Journal of Rational Mechanics and Analysis*, 4(6):323–425, 1955.
- R. S. Rivlin and D. W. Saunders. Large elastic deformations of isotropic materials VII. Experiments on the deformation of rubber. *Philosophical Transactions of the Royal Society of London A*, 243(865):251–288, 1951.
- G. Saccomandi. Universal solutions and relations in finite elasticity. In *Topics in finite elasticity*, pages 95–130. Springer, 2001a.
- G. Saccomandi. Universal solutions and relations in finite elasticity. In M. Hayes and G. Saccomandi, editors, *Topics in Finite Elasticity*, pages 95–130. Springer, 2001b.
- S. Sadik and A. Yavari. Nonlinear anisotropic viscoelasticity. *Journal of the Mechanics and Physics of Solids*, 182:105461, 2024.
- M. F. Shojaei and A. Yavari. Compatible-strain mixed finite element methods for incompressible nonlinear elasticity. *Journal of Computational Physics*, 361:247–279, 2018.
- M. Singh and A. C. Pipkin. Note on Ericksen’s problem. *Zeitschrift für angewandte Mathematik und Physik*, 16(5):706–709, 1965.
- A. J. M. Spencer. Part iii. Theory of invariants. *Continuum Physics*, 1:239–353, 1971.
- J. J. Sylvester. Sur l’équation en matrices $px = xq$. *CR Acad. Sci. Paris*, 99(2):67–71, 1884.
- E. B. Tadmor, R. E. Miller, and R. S. Elliott. *Continuum Mechanics and Thermodynamics: From Fundamental Concepts to Governing Equations*. Cambridge University Press, 2012.
- C. Truesdell. *The Elements of Continuum Mechanics*. Springer-Verlag, 1966.
- Z. Wesolowski and A. Seeger. On the screw dislocation in finite elasticity. In *Mechanics of Generalized Continua. Proceedings of the IUTAM Symposium on the Generalized Cosserat Continuum and the Continuum Theory of Dislocations with Applications (Ed. E. Kröner, Springer, Berlin, etc., 1968)*, pages 294–297. Springer, 1968.
- A. Yavari. Universal deformations in inhomogeneous isotropic nonlinear elastic solids. *Proceedings of the Royal Society A*, 477(2253):20210547, 2021a.

- A. Yavari. On Eshelby's inclusion problem in nonlinear anisotropic elasticity. *Journal of Micromechanics and Molecular Physics*, 6(01):2150002, 2021b.
- A. Yavari. Universal displacements in inextensible fiber-reinforced linear elastic solids. *Mathematics and Mechanics of Solids*, pages 1–17, 2023. doi: 10.1177/10812865231181924.
- A. Yavari. Universal deformations and inhomogeneities in isotropic Cauchy elasticity. *Proceedings of the Royal Society A*, 2024. URL <https://arxiv.org/abs/2404.06235>.
- A. Yavari and A. Goriely. Riemann–Cartan geometry of nonlinear dislocation mechanics. *Archive for Rational Mechanics and Analysis*, 205(1):59–118, 2012a.
- A. Yavari and A. Goriely. Weyl geometry and the nonlinear mechanics of distributed point defects. *Proceedings of the Royal Society A*, 468(2148):3902–3922, 2012b.
- A. Yavari and A. Goriely. Riemann–Cartan geometry of nonlinear disclination mechanics. *Mathematics and Mechanics of Solids*, 18(1):91–102, 2013a.
- A. Yavari and A. Goriely. Nonlinear elastic inclusions in isotropic solids. *Proc. R. Soc. A*, 469(2160):20130415, 2013b.
- A. Yavari and A. Goriely. The geometry of discombinations and its applications to semi-inverse problems in anelasticity. *Proceedings of the Royal Society A*, 470(2169):20140403, 2014.
- A. Yavari and A. Goriely. The twist-fit problem: Finite torsional and shear eigenstrains in nonlinear elastic solids. *Proceedings of the Royal Society A*, 471(2183):20150596, 2015.
- A. Yavari and A. Goriely. The anelastic Ericksen problem: Universal eigenstrains and deformations in compressible isotropic elastic solids. *Proc. R. Soc. A*, 472(2196):20160690, 2016.
- A. Yavari and A. Goriely. Universal deformations in anisotropic nonlinear elastic solids. *Journal of the Mechanics and Physics of Solids*, 156:104598, 2021.
- A. Yavari and A. Goriely. Universality in anisotropic linear anelasticity. *Journal of Elasticity*, 150(2):241–259, 2022a.
- A. Yavari and A. Goriely. The universal program of linear elasticity. *Mathematics and Mechanics of Solids*, 2022b.
- A. Yavari and A. Goriely. The universal program of nonlinear hyperelasticity. *Journal of Elasticity*, 154(1):91–146, 2023.
- A. Yavari and S. P. Pradhan. Accretion mechanics of nonlinear elastic circular cylindrical bars under finite torsion. *Journal of Elasticity*, 152(1-2):29–60, 2022.
- A. Yavari, C. Goodbrake, and A. Goriely. Universal displacements in linear elasticity. *Journal of the Mechanics and Physics of Solids*, 135:103782, 2020.
- A. Yavari, Y. Safa, and A. Soleiman Fallah. Finite extension of accreting nonlinear elastic solid circular cylinders. *Continuum Mechanics and Thermodynamics*, pages 1–17, 2023.
- L. M. Zubov. *Nonlinear Theory of Dislocations and Disclinations in Elastic Bodies*, volume 47. Springer Science & Business Media, 1997.